arXiv:1207.4757v1 [math.AC] 19 Jul 2012

Multivariate Difference-Differential Dimension Polynomials

Alexander Levin
Department of Mathematics, The Catholic University of America
Washington, D. C. 20064
E-mail: levin@cua.edu

Abstract

In this paper we generalize the Ritt-Kolchin method of characteristic sets and the classical Gröbner basis technique to prove the existence and obtain methods of computation of multivariate difference-differential dimension polynomials associated with a finitely generated difference-differential field extension. We also give an interpretation of such polynomials in the spirit of the A. Einstein's concept of strength of a system of PDEs and determine their invariants, that is, characteristics of a finitely generated difference-differential field extension carried by every its dimension polynomial.

1 Introduction

The role of Hilbert polynomials in commutative and homological algebra as well as in algebraic geometry and combinatorics is well known. A similar role in differential algebra is played by differential dimension polynomials, which describe in exact terms the freedom degree of a dynamic system as well as the number of arbitrary constants in the general solution of a system of partial algebraic differential equations.

The notion of differential dimension polynomial was introduced by E. Kolchin in 1964 [4] who proved the following fundamental result.

Theorem 1.1 *Let K be a differential field ($Char\, K = 0$), that is, a field considered together with the action of a set $\Delta = \{\delta_1, \dots, \delta_m\}$ of mutually commuting derivations of K into itself. Let Θ denote the free commutative semigroup of all power products of the form $\theta = \delta_1^{k_1} \dots \delta_m^{k_m}$ ($k_i \geq 0$), let $ord\, \theta = \sum_{i=1}^{m} k_i$, and for any $r \geq 0$, let $\Theta(r) = \{\theta \in \Theta \,|\, ord\, \theta \leq r\}$. Furthermore, let $L = K\langle \eta_1, \dots, \eta_n \rangle$ be a differential field extension of K generated by a finite set $\eta = \{\eta_1, \dots, \eta_n\}$. (As a field, $L = K(\{\theta\eta_j | \theta \in \Theta, 1 \leq j \leq n\})$.)*

Then there exists a polynomial $\omega_{\eta|K}(t) \in \mathbf{Q}[t]$ such that

(i) *$\omega_{\eta|K}(r) = trdeg_K K(\{\theta\eta_j | \theta \in \Theta(r), 1 \leq j \leq n\})$ for all sufficiently large $r \in \mathbf{Z}$;*

(ii) *$\deg \omega_{\eta|K} \leq m$ and $\omega_{\eta|K}(t)$ can be written as $\omega_{\eta|K}(t) = \sum_{i=0}^{m} a_i \binom{t+i}{i}$ where $a_0, \dots, a_m \in \mathbf{Z}$;*

(iii) *$d = \deg \omega_{\eta|K}$, a_m and a_d do not depend on the choice of the system of Δ-generators η of the extension L/K (clearly, $a_d \neq a_m$ iff $d < m$, that is $a_m = 0$). Moreover, a_m is equal to the differential transcendence degree of L*

over K, that is, to the maximal number of elements $\xi_1, \dots, \xi_k \in L$ such that the set $\{\theta\xi_i | \theta \in \Theta, 1 \leq i \leq k\}$ is algebraically independent over K.

In 1980 A. Mikhalev and E. Pankrat'ev [10] showed that a system of algebraic differential equations can be characterized by certain differential dimension polynomial, which expresses the strength of the system in the sense of A. Einstein. The concept of strength, which is an important characteristic of a system of PDEs governing a physical field, was described by A. Einstein as follows (see [2]): "... the system of equations is to be chosen so that the field quantities are determined as strongly as possible. In order to apply this principle, we propose a method which gives a measure of strength of an equation system. We expand the field variables, in the neighborhood of a point $\mathcal{P}$, into a Taylor series (which presupposes the analytic character of the field); the coefficients of these series, which are the derivatives of the field variables at $\mathcal{P}$, fall into sets according to the degree of differentiation. In every such degree there appear, for the first time, a set of coefficients which would be free for arbitrary choice if it were not that the field must satisfy a system of differential equations. Through this system of differential equations (and its derivatives with respect to the coordinates) the number of coefficients is restricted, so that in each degree a smaller number of coefficients is left free for arbitrary choice. The set of numbers of "free" coefficients for all degrees of differentiation is then a measure of the "weakness" of the system of equations, and through this, also of its "strength"."

In this paper we generalize the Ritt-Kolchin method of characteristic sets to the case of difference-differential polynomials and apply this method to prove the existence and find invariants of multivariate dimension polynomials associated with a fixed partition of the basic sets of derivations and automorphisms. We show that one can assign such a multivariate polynomial to a system of partial algebraic difference-differential equations, and this polynomial expresses the strength of the system in the sense of A. Einstein. We also find new invariants of a finitely generated difference-differential field extension carried by multivariate dimension polynomials, that is, characteristics of the extension, which do not depend on the choice of the system of its generators.

2 Preliminaries

In this section we present some basic concepts and results used in the rest of the paper.

Throughout the paper, $\mathbf{N}, \mathbf{Z}$, $\mathbf{Q}$, and $\mathbf{R}$ denote the sets of all non-negative integers, integers, rational numbers, and real numbers, respectively. As usual, $\mathbf{Q}[t]$ denotes the ring of polynomials in one variable t with rational coefficients. By a ring we always mean an associative ring with a unity. Every ring homomorphism is unitary (maps unity onto unity), every subring of a ring contains the unity of the ring. Unless otherwise indicated, by a module over a ring R we always mean a unitary left R-module.

By a *difference-differential ring* we mean a commutative ring R together with finite sets $\Delta = \{\delta_1, \dots, \delta_m\}$ and $\sigma = \{\alpha_1, \dots, \alpha_n\}$ of derivations and automorphisms of R, respectively, such that any two mappings of the set $\Delta \bigcup \sigma$ commute. The set $\Delta \bigcup \sigma$ is called the *basic set* of the difference-differential ring R, which is also called a $\Delta \bigcup \sigma$-ring. If R is a field, it is called a *difference-differential field* or a $\Delta \bigcup \sigma$-field. Furthermore, in what follows, we denote the set $\{\alpha_1, \dots, \alpha_n, \alpha_1^{-1}, \dots, \alpha_n^{-1}\}$ by σ^*.

If R is a difference-differential ring with a basic set $\Delta \bigcup \sigma$ described above, then Λ will denote the free commutative semigroup of all power products of the form $\lambda = \delta_1^{k_1} \dots \delta_m^{k_m} \alpha_1^{l_1} \dots \alpha_n^{l_n}$ where $k_i \in \mathbf{N}$, $l_j \in \mathbf{Z}$ $(1 \leq i \leq m,\ 1 \leq j \leq n)$. For any such an element λ, we set $\lambda_\Delta = \delta_1^{k_1} \dots \delta_m^{k_m}$, $\lambda_\sigma = \alpha_1^{l_1} \dots \alpha_n^{l_n}$, and denote by Λ_Δ and Λ_σ the commutative semigroup of power products $\delta_1^{k_1} \dots \delta_m^{k_m}$ and the commutative group of elements of the form $\alpha_1^{l_1} \dots \alpha_n^{l_n}$, respectively. The *order* of λ is defined as $ord\, \lambda = \sum_{i=1}^{m} k_i + \sum_{j=1}^{n} |l_j|$, and for every $r \in \mathbf{N}$, we set $\Lambda(r) = \{\lambda \in \Lambda \,|\, ord\, \lambda \leq r\}$ $(r \in \mathbf{N})$.

A subring (ideal) R_0 of a Δ-σ-ring R is said to be a difference-differential (or Δ-σ-) subring of R (respectively, difference-differential (or Δ-σ-) ideal of R) if R_0 is closed with respect to the action of any operator of $\Delta \bigcup \sigma^*$. In this case the restriction of a mapping from $\Delta \bigcup \sigma^*$ on R_0 is denoted by same symbol. If a prime (maximal) ideal P of R is closed with respect to the action of $\Delta \bigcup \sigma^*$, it is called a *prime* (respectively, *maximal*) difference-differential (or Δ-σ-) *ideal* of R.

If R is a Δ-σ-field and R_0 a subfield of R which is also a Δ-σ-subring of R, then R_0 is said to be a Δ-σ-subfield of R; R, in turn, is called a difference-differential (or Δ-σ-) field extension or a Δ-σ-overfield of R_0. In this case we also say that we have a Δ-σ-field extension R/R_0.

If R is a Δ-σ-ring and $\Sigma \subseteq R$, then the intersection of all Δ-σ-ideals of R containing the set Σ is, obviously, the smallest Δ-σ-ideal of R containing Σ. This ideal is denoted by $[\Sigma]$. (It is clear that $[\Sigma]$ is generated, as an ideal, by the set $\{\lambda\xi | \xi \in \Sigma,\ \lambda \in \Lambda\}$). If the set Σ is finite, $\Sigma = \{\xi_1, \dots, \xi_q\}$, we say that the Δ-σ-ideal $I = [\Sigma]$ is finitely generated (we write this as $I = [\xi_1, \dots, \xi_q]$) and call $\xi_1, \dots, \xi_q$ differential or Δ-σ-generators of I.

If K_0 is a Δ-σ-subfield of a Δ-σ-field K and $\Sigma \subseteq K$, then the intersection of all Δ-σ-subfields of K containing K_0 and Σ is the unique Δ-σ-subfield of K containing K_0 and Σ and contained in every Δ-σ-subfield of K containing K_0 and Σ. It is denoted by $K_0\langle\Sigma\rangle$. If $K = K_0\langle\Sigma\rangle$ and the set Σ is finite, $\Sigma = \{\eta_1, \dots, \eta_s\}$, then K is said to be a finitely generated Δ-σ-extension of K_0 with the set of Δ-σ-generators $\{\eta_1, \dots, \eta_s\}$. In this case we write $K = K_0\langle\eta_1, \dots, \eta_s\rangle$. It is easy to see that the field $K_0\langle\eta_1, \dots, \eta_s\rangle$ coincides with the field $K_0(\{\lambda\eta_i | \lambda \in \Lambda, 1 \leq i \leq s\})$.

Let R and S be two difference-differential rings with the same basic set $\Delta \bigcup \sigma$, so that elements of the sets Δ and σ act on each of the rings as mutually commuting derivations and automorphisms, respectively, and every two mapping of the set $\Delta \bigcup \sigma$ commute. (More rigorously, we assume that there exist injective mappings of the sets Δ and σ into the sets of derivations and automorphisms of the rings R and S, respectively, such that the images of any two elements of $\Delta\bigcup\sigma$ commute. For convenience we will denote the images of elements of $\Delta\bigcup\sigma$ under these mappings by the same symbols $\delta_1,\dots,\delta_m,\alpha_1,\dots,\alpha_n$). A ring homomorphism $\phi : R \longrightarrow S$ is called a *difference-differential* or Δ-σ-*homomorphism* if $\phi(\tau a) = \tau\phi(a)$ for any $\tau \in \Delta\bigcup\sigma$, $a \in R$. The notions of Δ-σ-*epimorphism,* Δ-σ-*monomorphism,* Δ-σ-*automorphism*, etc. are defined naturally (as the corresponding ring homomorphisms that are Δ-σ-homomorphisms).

If K is a difference-differential (Δ-σ-) field and $Y = \{y_1,\dots,y_s\}$ is a finite set of symbols, then one can consider the countable set of symbols $\Lambda Y = \{\lambda y_j|\lambda \in \Lambda, 1 \leq j \leq s\}$ and the polynomial ring $R = K[\{\lambda y_j|\lambda \in \Lambda, 1 \leq j \leq s\}]$ in the set of indeterminates ΛY over the field K. This polynomial ring is naturally viewed as a Δ-σ-ring where $\tau(\lambda y_j) = (\tau\lambda)y_j$ for any $\tau \in \Delta\bigcup\sigma$, $\lambda \in \Lambda$, $1 \leq j \leq s$, and the elements of Δ act on the coefficients of the polynomials of R as they act in the field K. The ring R is called a *ring of difference-differential* (or Δ-σ-) *polynomials* in the set of differential (Δ-σ-) indeterminates $y_1,\dots,y_s$ over K. This ring is denoted by $K\{y_1,\dots,y_s\}$ and its elements are called difference-differential (or Δ-σ-) polynomials.

Let $L = K\langle\eta_1,\dots,\eta_s\rangle$ be a difference-differential field extension of K generated by a finite set $\eta = \{\eta_1,\dots,\eta_s\}$.

The following is a unified version of E. Kolchin's theorem on differential dimension polynomial and the author's theorem on the dimension polynomial of a difference field extension (see [7] or [Levin(2008), Theorem 4.2.5]).

Theorem 2.1 *With the above notation, there exists a polynomial* $\phi_{\eta|K}(t) \in \mathbf{Q}[t]$ *such that*

(i) $\phi_{\eta|K}(r) = trdeg_K K(\{\lambda\eta_j|\lambda \in \Lambda(r), 1 \leq j \leq s\})$ *for all sufficiently large* $r \in \mathbf{Z}$*;*

(ii) $\deg\phi_{\eta|K} \leq m+n$ *and* $\phi_{\eta|K}(t)$ *can be written as* $\phi_{\eta|K}(t) = \sum_{i=0}^{m+n} a_i \binom{t+i}{i}$ *where* $a_0,\dots,a_{m+n} \in \mathbf{Z}$ *and* $2^n|a_{m+n}$.

(iii) $d = \deg\phi_{\eta|K}$*,* a_{m+n} *and* a_d *do not depend on the set of difference-differential generators* η *of* L/K *(*$a_d \neq a_{m+n}$ *if and only if* $d < m+n$*). Moreover,* $\frac{a_{m+n}}{2^n}$ *is equal to the difference-differential transcendence degree of* L *over* K *(denoted by* Δ-σ-$trdeg_K L$*), that is, to the maximal number of elements* $\xi_1,\dots,\xi_k \in L$ *such that the family* $\{\lambda\xi_i|\lambda \in \Lambda, 1 \leq i \leq k\}$ *is algebraically independent over* K*.*

The polynomial whose existence is established by this theorem is called a *difference-differential* (or Δ-σ-) *dimension polynomial* of the extension L/K associated with the system of difference-differential generators η.

3 Partitions of the basic set of derivations and the formulation of the main theorem

Let K be a difference-differential field of zero characteristic with basic sets $\Delta = \{\delta_1, \dots, \delta_m\}$ and $\sigma = \{\alpha_1, \dots, \alpha_n\}$ of derivations and automorphisms, respectively. Suppose that the set of derivations is represented as the union of p disjoint subsets ($p \geq 1$):

$$\Delta = \Delta_1 \bigcup \dots \bigcup \Delta_p \tag{3. 1}$$

where

$$\Delta_1 = \{\delta_1, \dots, \delta_{m_1}\},\ \Delta_2 = \{\delta_{m_1+1}, \dots, \delta_{m_1+m_2}\},\ \dots,$$

$$\Delta_p = \{\delta_{m_1+\dots+m_{p-1}+1}, \dots, \delta_m\}.\ (m_1 + \dots + m_p = m).$$

In other words, we fix a partition of the set Δ into p sets of derivations.

If $\lambda = \delta_1^{k_1} \dots \delta_m^{k_m} \alpha_1^{l_1} \dots \alpha_n^{l_n} \in \Lambda$ ($k_i \in \mathbf{N},\ l_j \in \mathbf{Z}$), then the order of λ with respect to a set Δ_i ($1 \leq i \leq p$) is defined as $\sum\limits_{\nu=m_1+\dots+m_{i-1}+1}^{m_1+\dots+m_i} k_\nu$; it is denoted by $ord_i\lambda$. (If $i = 1$, the last sum is replaced by $k_1 + \dots + k_{m_1}$.) The number $ord_\sigma\lambda = \sum\limits_{j=1}^{n} |l_j|$ is called the order of λ with respect to σ.

If $r_1, \dots, r_{p+1} \in \mathbf{N}$, we set

$\Lambda(r_1, \dots, r_{p+1}) = \{\lambda \in \Lambda \,|\, ord_i\lambda \leq r_i \text{ for } i = 1, \dots, p \text{ and } ord_\sigma\lambda \leq r_{p+1}\}$.

In what follows, for any permutation $(j_1, \dots, j_{p+1})$ of the set $\{1, \dots, p+1\}$, $<_{j_1,\dots,j_{p+1}}$ will denote the lexicographic order on $\mathbf{N}^{p+1}$ such that $(r_1, \dots, r_{p+1}) <_{j_1,\dots,j_{p+1}} (s_1, \dots, s_{p+1})$ if and only if either $r_{j_1} < s_{j_1}$ or there exists $k \in \mathbf{N}$, $1 \leq k \leq p$, such that $r_{j_\nu} = s_{j_\nu}$ for $\nu = 1, \dots, k$ and $r_{j_{k+1}} < s_{j_{k+1}}$. Furthermore, if $\Sigma \subseteq \mathbf{N}^{p+1}$, then Σ' denotes the set

$\{e \in \Sigma | e$ is a maximal element of Σ with respect to one of the $(p+1)!$ lexicographic orders $<_{j_1,\dots,j_{p+1}}\}$. For example, if $\Sigma = \{(3,0,2), (2,1,1), (0,1,4), (1,0,3), (1,1,6), (3,1,0), (1,2,0)\} \subseteq \mathbf{N}^3$, then $\Sigma' = \{(3,0,2), (3,1,0), (1,1,6), (1,2,0)\}$.

Theorem 3.1 *Let $L = K\langle\eta_1, \dots, \eta_s\rangle$ be a Δ-σ-field extension generated by a set $\eta = \{\eta_1, \dots, \eta_s\}$. Then there exists a polynomial $\Phi_\eta(t_1, \dots, t_{p+1})$ in $(p+1)$ variables $t_1, \dots, t_{p+1}$ with rational coefficients such that*

(i) $\Phi_\eta(r_1, \dots, r_{p+q}) = trdeg_K K(\bigcup\limits_{j=1}^{s} \Lambda(r_1, \dots, r_{p+1})\eta_j)$

for all sufficiently large $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$ (i. e., there exist $s_1, \dots, s_{p+1} \in \mathbf{N}$ such that the last equality holds for all $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$ with $r_1 \geq s_1, \dots, r_{p+1} \geq s_{p+1}$);

(ii) *$deg_{t_i}\Phi_\eta \le m_i$ $(1 \le i \le p)$ and $deg_{t_{p+1}}\Phi_\eta \le n$, so that $deg\,\Phi_\eta \le m+n$ and $\Phi_\eta(t_1,\dots,t_{p+1})$ can be represented as*

$$\Phi_\eta(t_1,\dots,t_{p+1}) = \sum_{i_1=0}^{m_1}\dots\sum_{i_p=0}^{m_p}\sum_{i_{p+1}=0}^{n} a_{i_1\dots i_{p+1}}\binom{t_1+i_1}{i_1}\dots\binom{t_{p+1}+i_{p+1}}{i_{p+1}}$$

where $a_{i_1\dots i_{p+1}} \in \mathbf{Z}$ and $2^n \mid a_{m_1\dots m_p n}$.

(iii) *Let $E_\eta = \{(i_1,\dots,i_{p+1}) \in \mathbf{N}^{p+1} \,|\, 0 \le i_k \le m_k$ for $k = 1,\dots,p$, $0 \le i_{p+1} \le n$, and $a_{i_1\dots i_{p+1}} \ne 0\}$. Then $d = deg\,\Phi_\eta$, $a_{m_1\dots m_p n}$, elements $(k_1,\dots,k_{p+1}) \in E'_\eta$, the corresponding coefficients $a_{k_1\dots k_{p+1}}$ and the coefficients of the terms of total degree d do not depend on the choice of the system of Δ-σ-generators η. Furthermore, $\dfrac{a_{m_1\dots m_p n}}{2^n} = \Delta$-$\sigma$-$tr.deg_K L$.*

Definition 3.2 *The polynomial $\Phi_\eta(t_1,\dots,t_{p+1})$ whose existence is established by Theorem 3.1 is called the difference-differential* (or Δ-σ-) *dimension polynomial associated with the partition (3.1) of the basic set of derivations.*

The Δ-σ-dimension polynomial associated with partition (3.1) has the following interpretation as the strength of a system of difference-differential equations.

Let us consider a system of partial difference-differential equations

$$A_i(f_1,\dots,f_s) = 0 \qquad (i = 1,\dots,q) \tag{3. 2}$$

over a field of of functions in m real variables $x_1,\dots,x_m$ ($f_1,\dots,f_s$ are unknown functions of $x_1,\dots,x_m$). Suppose that $\Delta = \{\delta_1,\dots,\delta_m\}$ where δ_i is the partial differentiation $\partial/\partial x_i$ $(i = 1,\dots,m)$ and the basic set of automorphisms $\sigma = \{\alpha_1,\dots,\alpha_m\}$ consists of m shifts of arguments, $f(x_1,\dots,x_m) \mapsto f(x_1,\dots,x_{i-1},x_i+h_i,x_{i+1},\dots,x_m)$ $(1 \le i \le m$, $h_1,\dots,h_m$ are some real numbers). Thus, we assume that the the left-hand sides of the equations in (3.2) contain unknown functions f_i, their partial derivatives, their images under the shifts α_j and various compositions of such shifts and partial derivations. Furthermore, we suppose that system (3.2) is algebraic, that is, all $A_i(y_1,\dots,y_s)$ are elements of a ring of Δ-σ-polynomials $K\{y_1,\dots,y_s\}$ with coefficients in some functional Δ-σ-field K.

Let us consider a grid with equal cells of dimension $h_1 \times \dots \times h_m$ that fills the whole space $\mathbf{R}^m$. Let us fix some node $\mathcal{P}$ say that *a node $\mathcal{Q}$ has order i* (with respect to $\mathcal{P}$) if the shortest path from $\mathcal{P}$ to $\mathcal{Q}$ along the edges of the grid consists of i steps (by a step we mean a path from a node of the grid to a neighbor node along the edge between these two nodes). Say, the orders of the nodes in the two-dimensional case are as follows (a number near a node shows the order of this node).



5	4	3	2	3	4	5
4	3	2	1	2	3	4
3	2	1	$\mathcal{P}$	1	2	3
4	3	2	1	2	3	4
5	4	3	2	3	4	5

Let us fix partition (3.1) of the set of basic derivations Δ (such a partition can be, for example, a natural separation of (all or some) derivations with respect to coordinates and the derivation with respect to time). Let us consider the values of the unknown functions $f_1, \dots, f_s$ and their partial derivatives whose ith order (that is the order with respect to the derivations of the set Δ_i) does not exceed r_i $(1 \leq i \leq p)$ at the nodes whose order does not exceed r_{p+1} $(r_1, \dots, r_{p+1} \in \mathbf{N})$. If $f_1, \dots, f_s$ should not satisfy any system of equations (or any other condition), these values can be chosen arbitrarily. Because of the system (and equations obtained from the equations of the system by partial differentiations and transformations of the form $f_j(x_1, \dots, x_m) \mapsto f_j(x_1 + k_1h_1, \dots, x_m + k_mh_m)$ with $k_1, \dots, k_m \in \mathbf{Z}$, $1 \leq j \leq s$), the number of independent values of the functions $f_1, \dots, f_s$ and their partial derivatives whose ith order does not exceed r_i $(1 \leq i \leq p)$ at the nodes of order $\leq r_{p+1}$ decreases. This number, which is a function of $p+1$ variables $r_1, \dots, r_{p+1}$, is the "measure of strength" of the system in finite differences in the sense of A. Einstein). We denote it by $S_{r_1, \dots, r_{p+1}}$.

Suppose that the Δ-σ-ideal J generated in $K\{y_1, \dots, y_s\}$ by the Δ-σ-polynomials $A_1, \dots, A_q$ is prime (e. g., the polynomials are linear). The we say that the system of difference-differential equations (3.2) is prime. In this case, the field of fractions L of the Δ-σ-integral domain $K\{y_1, \dots, y_s\}/J$ has a natural structure of a Δ-σ-field extension of K generated by the finite set $\eta = \{\eta_1, \dots, \eta_s\}$ where η_i is the canonical image of y_i in $K\{y_1, \dots, y_s\}/J$ $(1 \leq i \leq s)$. It is easy to see that the Δ-σ-dimension polynomial $\Phi_\eta(t_1, \dots, t_{p+1})$ of the extension L/K associated with the system of Δ-σ-generators η has the property that $\Phi_\eta(r_1, \dots, r_{p+1}) = S_{r_1, \dots, r_{p+1}}$ for all sufficiently large $(r_1, \dots, r_{p+q}) \in \mathbf{N}^{p+1}$, so this dimension polynomial is the measure of strength of the system of difference-differential equations (3.2) in the sense of A. Einstein.

4 Numerical polynomials of subsets of $\mathbf{N}^m \times \mathbf{Z}^n$

Definition 4.1 *A polynomial $f(t_1, \dots, t_p)$ in p variables $t_1, \dots, t_p$ $(p \in \mathbf{N}, p \geq 1)$ with rational coefficients is called* numerical *if $f(t_1, \dots, t_p) \in \mathbf{Z}$ for all sufficiently large $(t_1, \dots, t_p) \in \mathbf{Z}^p$. (Recall that it means that there exist integers*

$s_1, \dots, s_p$ such that $f(r_1, \dots, r_p) \in \mathbf{Z}$ as soon as $(r_1, \dots, r_p) \in \mathbf{Z}^p$ and $r_i \geq s_i$ for all $i = 1, \dots, p$.)

It is clear that every polynomial with integer coefficients is numerical. As an example of a numerical polynomial in p variables with noninteger coefficients ($p \in \mathbf{N}, p \geq 1$) one can consider a polynomial $\prod_{i=1}^{p} \binom{t_i}{m_i}$ where $m_1, \dots, m_p \in \mathbf{N}$. (As usual, $\binom{t}{k}$ ($k \in \mathbf{Z}, k \geq 1$) denotes the polynomial $\frac{t(t-1)\dots(t-k+1)}{k!}$ in one variable t, $\binom{t}{0} = 1$, and $\binom{t}{k} = 0$ if k is a negative integer.)

The following theorem proved in [6] gives the "canonical" representation of a numerical polynomial in several variables.

Theorem 4.2 *Let $f(t_1, \dots, t_p)$ be a numerical polynomial in p variables $t_1, \dots, t_p$, and let $deg_{t_i} f = m_i$ ($m_1, \dots, m_p \in \mathbf{N}$). Then the polynomial $f(t_1, \dots, t_p)$ can be represented in the form*

$$f(t_1, \dots t_p) = \sum_{i_1=0}^{m_1} \dots \sum_{i_p=0}^{m_p} a_{i_1 \dots i_p} \binom{t_1 + i_1}{i_1} \dots \binom{t_p + i_p}{i_p} \tag{4. 1}$$

with integer coefficients $a_{i_1 \dots i_p}$ ($0 \leq i_k \leq m_k$ for $k = 1, \dots, p$) that are uniquely defined by the numerical polynomial.

In what follows (until the end of the section), we deal with subsets of the set $\mathbf{N}^m \times \mathbf{Z}^n$ (m and n are positive integers). Furthermore, we fix a partition of the set $\mathbf{N}_m = \{1, \dots, m\}$ into p disjoint subsets ($p \geq 1$):

$$\mathbf{N}_m = \{1, \dots, m_1\} \bigcup \{m_1+1, \dots, m_1+m_2\} \bigcup \dots \bigcup \{m_1+\dots+m_{p-1}+1, \dots, m\} \tag{4. 2}$$

($m_1 + \dots + m_p = m$).

If $a = (a_1, \dots, a_{m+n}) \in \mathbf{N}^m \times \mathbf{Z}^n$ we denote the numbers $\sum_{i=1}^{m_1} a_i$, $\sum_{i=m_1+1}^{m_1+m_2} a_i, \dots,$ $\sum_{i=m_1+\dots+m_{p-1}+1}^{m} a_i$, $\sum_{i=m+1}^{m+n} |a_i|$ by $ord_1 a, \dots, ord_{p+1} a$, respectively.

As in [6, Section 2.5], let us consider the set $\mathbf{Z}^n$ as a union

$$\mathbf{Z}^n = \bigcup_{1 \leq j \leq 2^n} \mathbf{Z}_j^{(n)} \tag{4. 3}$$

where $\mathbf{Z}_1^{(n)}, \dots, \mathbf{Z}_{2^n}^{(n)}$ are all different Cartesian products of n sets each of which is either $\mathbf{N}$ or $\mathbf{Z}_- = \{k \in \mathbf{Z} \,|\, k \leq 0\}$. We assume that $\mathbf{Z}_1^{(n)} = \mathbf{N}^n$ and call $\mathbf{Z}_j^{(n)}$ the *jth orthant* of the set $\mathbf{Z}^n$ ($1 \leq j \leq 2^n$). Furthermore, we consider $\mathbf{N}^m \times \mathbf{Z}^n$ as a partially ordered set with the order $\trianglelefteq$ such that $(e_1, \dots, e_m, f_1, \dots, f_n) \trianglelefteq$

$(e_1', \dots, e_m', f_1', \dots, f_n')$ if and only if $(f_1, \dots, f_n)$ and $(f_1', \dots, f_n')$ belong to the same orthant $\mathbf{Z}_k^{(n)}$ $(1 \leq k \leq 2^n)$ and the $(m+n)$-tuple $(e_1, \dots, e_m, |f_1|, \dots, |f_n|)$ is less than $(e_1', \dots, e_m', |f_1'|, \dots, |f_n'|)$ with respect to the product order on $\mathbf{N}^{m+n}$.

In what follows, for any set $A \subseteq \mathbf{N}^m \times \mathbf{Z}^n$, W_A will denote the set of all elements of $\mathbf{N}^m \times \mathbf{Z}^n$ that do not exceed any element of A with respect to the order $\trianglelefteq$. (Thus, $w \in W_A$ if and only if there is no element $a \in A$ such that $a \trianglelefteq w$.) Furthermore, for any $r_1, \dots r_{p+1} \in \mathbf{N}$, $A(r_1, \dots r_{p+1})$ will denote the set of all elements $x = (x_1, \dots, x_m, x_1', \dots, x_n') \in A$ such that $ord_i x \leq r_i$ $(i = 1, \dots, p+1)$.

The above notation can be naturally restricted to subsets of $\mathbf{N}^m$. If $E \subseteq \mathbf{N}^m$ and $s_1, \dots, s_p$ are nonnegative integers, then $E(s_1, \dots, s_p)$ will denote the set of all m-tuples $e = (e_1, \dots, e_m) \in E$ such that $ord_i(e_1, \dots, e_m, 0) \leq s_i$ for $i = 1, \dots, p$. Furthermore, we shall associate with a set $E \subseteq \mathbf{N}^m$ a set $V_E \subseteq \mathbf{N}^m$ that consists of all m-tuples $v = (v_1, \dots, v_m) \in \mathbf{N}$ that are not greater than or equal to any m-tuple in E with respect to the product order on $\mathbf{N}^m$. (Recall that the product order on $\mathbf{N}^m$ is a partial order $\leq_P$ on $\mathbf{N}^m$ such that $c = (c_1, \dots, c_m) \leq_P c' = (c_1', \dots, c')$ if and only if $c_i \leq c_i'$ for all $i = 1, \dots, m$. If $c \leq_P c'$ and $c \neq c'$, we write $c <_P c'$). Clearly, an element $v = (v_1, \dots, v_m) \in \mathbf{N}^m$ belongs to V_E if and only if for any element $(e_1, \dots, e_m) \in E$ there exists $i \in \mathbf{N}, 1 \leq i \leq m$, such that $e_i > v_i$.

The following two theorems proved in [6, Chapter 2] generalize the well-known Kolchin's result on the numerical polynomials associated with subsets of $\mathbf{N}$ (see [5, Chapter 0, Lemma 17]) and give the explicit formula for the numerical polynomials in p variables associated with a finite subset of $\mathbf{N}^m$.

Theorem 4.3 *Let E be a subset of $\mathbf{N}^m$ where $m = m_1 + \dots + m_p$ for some nonnegative integers $m_1, \dots, m_p$ $(p \geq 1)$. Then there exists a numerical polynomial $\omega_E(t_1, \dots, t_p)$ with the following properties:*

(i) *$\omega_E(r_1, \dots, r_p) = Card\, V_E(r_1, \dots, r_p)$ for all sufficiently large $(r_1, \dots, r_p) \in \mathbf{N}^p$*

(as usual, $Card\, M$ denotes the number of elements of a finite set M).

(ii) *The total degree of the polynomial ω_E does not exceed m and $deg_{t_i} \omega_E \leq m_i$ for all $i = 1, \dots, p$.*

(iii) *$deg\, \omega_E = m$ if and only if the $E = \emptyset$. Then $\omega_E(t_1, \dots, t_p) = \prod_{i=1}^{p} \binom{t_i + m_i}{m_i}$.*

(iv) *ω_E is a zero polynomial if and only if $(0, \dots, 0) \in E$.*

Definition 4.4 *The polynomial $\omega_E(t_1, \dots, t_p)$ whose existence is stated by Theorem 4.3 is called the dimension polynomial of the set $E \subseteq \mathbf{N}^m$ associated with the partition $(m_1, \dots, m_p)$ of m. If $p = 1$, the polynomial ω_E is called the Kolchin polynomial of the set E.*

Theorem 4.5 *Let $E = \{e_1, \dots, e_q\}$ be a finite subset of $\mathbf{N}^m$ where q is a positive integer and $m = m_1 + \dots + m_p$ for some nonnegative integers $m_1, \dots, m_p$*

($p \geq 1$). Let $e_i = (e_{i1}, \dots, e_{im})$ ($1 \leq i \leq q$) and for any $l \in \mathbf{N}$, $0 \leq l \leq q$, let $\Gamma(l, q)$ denote the set of all l-element subsets of the set $\mathbf{N}_q = \{1, \dots, q\}$. Furthermore, for any $\sigma \in \Gamma(l, q)$, let $\bar{e}_{\sigma j} = \max\{e_{ij} | i \in \sigma\}$ ($1 \leq j \leq m$) and $b_{\sigma j} = \sum\limits_{h \in \sigma j} \bar{e}_{\sigma h}$. Then

$$\omega_E(t_1, \dots, t_p) = \sum_{l=0}^{q} (-1)^l \sum_{\sigma \in \Gamma(l,q)} \prod_{j=1}^{p} \binom{t_j + m_j - b_{\sigma j}}{m_j} \qquad (4.\ 4)$$

Remark. It is clear that if E is any subset of $\mathbf{N}^m$ and E^* is the set of all minimal elements of the set E with respect to the product order on $\mathbf{N}^m$, then the set E^* is finite and $\omega_E(t_1, \dots, t_p) = \omega_{E^*}(t_1, \dots, t_p)$. Thus, Theorem 4.3 gives an algorithm that allows one to find a numerical polynomial associated with any subset of $\mathbf{N}^m$ (and with a given partition of the set $\{1, \dots, m\}$): one should first find the set of all minimal points of the subset and then apply Theorem 4.3.

The following result can be obtained precisely in the same way as Theorem 3.4 of [8] (the only difference is that the proof in the mentioned paper uses Theorem 3.2 of [8] in the case $p = 2$, while the proof of the theorem below should refer to the Theorem 3.2 of [8] where p is any positive integer).

Theorem 4.6 *Let A be a subset of $\mathbf{N}^m \times \mathbf{Z}^n$ and let partition (4.2) of the set $\mathbf{N}_m$ be fixed. Then there exists a numerical polynomial $\phi_A(t_1, \dots, t_{p+1})$ in $p+1$ variables $t_1, \dots, t_{p+1}$ with the following properties.*

(i) *$\phi_A(r_1, \dots, r_{p+1}) = Card\, W_A(r_1, \dots, r_{p+1})$ for all sufficiently large $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$.*

(ii) *$deg_{t_i} \phi_A \leq m_i$ for $i = 1, \dots, p$ and $deg_{t_{p+1}} \phi_A \leq n$.*

(iii) *Let us consider a mapping $\rho : \mathbf{N}^m \times \mathbf{Z}^n \longrightarrow \mathbf{N}^{m+2n}$ such that*

$\rho((a_1, \dots, a_{m+n}) = (a_1, \dots, a_m, \max\{a_{m+1}, 0\}, \dots, \max\{-a_{m+1}, 0\},$
$\max\{a_{m+n}, 0\}, \dots, \max\{-a_{m+n}, 0\})$.

Let $B = \rho(A) \bigcup \{e_1, \dots, e_n\}$ where e_i ($1 \leq i \leq n$) is a $(m + 2n)$-tuple in $\mathbf{N}^{m+2n}$ whose $(m + i)$th and $(m + n + i)$th coordinates are equal to 1 and all other coordinates are equal to 0. Then $\phi_A(t_1, \dots, t_{p+1}) = \omega_B(t_1, \dots, t_{p+1})$ where $\omega_B(t_1, \dots, t_{p+1})$ is the dimension polynomial of the set B (see Definition *4.4) associated with the partition $\mathbf{N}_{m+2n} = \{1, \dots, m_1\} \bigcup \{m_1 + 1, \dots, m_1 + m_2\} \bigcup \dots \bigcup \{m_1 + \dots + m_{p-1} + 1, \dots, m\} \bigcup \{m+1, \dots, m+2n\}$ of the set $\mathbf{N}_{m+2n}$.*

(iv) *If $A = \emptyset$, then*

$$\phi_A(t_1, \dots, t_{p+1}) = \binom{t_1 + m_1}{m_1} \dots \binom{t_p + m_p}{m_p} \sum_{i=0}^{n} (-1)^{n-i} 2^i \binom{n}{i} \binom{t_{p+1} + i}{i}.$$

(v) *$\phi_A(t_1, \dots, t_{p+1}) = 0$ if and only if $(0, \dots, 0) \in A$.*

5 Proof of the main theorem and computation of difference-differential dimension polynomials via characteristic sets

In this and next sections we prove Theorem 3.1 and give two methods of computation of difference-differential dimension polynomials of Δ-σ-field extensions. One of them is based on constructing a characteristic set of the defining prime Δ-σ-ideal of the extension; the other approach is the computation of the dimension polynomial of the associated module of Käller differentials via (generalized) Gröbner basis method. Both approaches use total term orderings with respect to several orders defined by partitions of the basic sets of derivations and automorphisms.

In what follows we use the notation and conventions introduced at the beginning of section 3. In particular, we assume that partition (3.1) of the set of basic derivations $\Delta = \{\delta_1, \dots, \delta_m\}$ is fixed.

Let us consider $p+1$ total orderings $<_1, \dots, <_p, <_\sigma$ of the set of power products Λ such that

$\lambda = \delta_1^{k_1} \dots \delta_m^{k_m} \alpha_1^{l_1} \dots \alpha_n^{l_n} <_i \lambda' = \delta_1^{k'_1} \dots \delta_m^{k'_m} \alpha_1^{l'_1} \dots \alpha_n^{l'_n}$ $(1 \leq i \leq p)$ if and only if $(ord_i\lambda, ord\,\lambda, ord_1\lambda, \dots, ord_{i-1}\lambda, ord_{i+1}\lambda, \dots, ord_p\lambda, ord_\sigma\lambda, k_{m_1+\dots+m_{i-1}+1}, \dots, k_{m_1+\dots+m_i}, k_1, \dots, k_{m_1+\dots+m_{i-1}}, k_{m_1+\dots+m_i+1}, \dots, k_m, |l_1|, \dots, |l_n|, l_1, \dots, l_n)$ is less than $(ord_i\lambda', ord\,\lambda', ord_1\lambda', \dots, ord_{i-1}\lambda', ord_{i+1}\lambda', \dots, ord_p\lambda', ord_\sigma\lambda', k'_{m_1+\dots+m_{i-1}+1}, \dots, k'_{m_1+\dots+m_i}, k'_1, \dots, k'_{m_1+\dots+m_{i-1}}, k'_{m_1+\dots+m_i+1}, \dots, k'_m, |l'_1|, \dots, |l'_n|, l'_1, \dots, l'_n)$ with respect to the lexicographic order on $\mathbf{N}^{m+2n+p+2}$.

Similarly, $\lambda <_\sigma \lambda'$ if and only if $(ord_\sigma\lambda, ord\,\lambda, ord_1\lambda, \dots, ord_p\lambda, |l_1|, \dots, |l_n|, l_1, \dots, l_n, k_1, \dots, k_m)$ is less than the corresponding $(m+2n+p+2)$-tuple for λ' with respect to the lexicographic order on $\mathbf{N}^{m+2n+p+2}$.

Two elements $\lambda_1 = \delta_1^{k_1} \dots \delta_m^{k_m} \alpha_1^{l_1} \dots \alpha_n^{l_n}$ and $\lambda_2 = \delta_1^{r_1} \dots \delta_m^{r_m} \alpha_1^{s_1} \dots \alpha_n^{s_n}$ in Λ are called *similar*, if the n-tuples $(l_1, \dots, l_n)$ and $(s_1, \dots, s_n)$ belong to the same orthant of $\mathbf{Z}^n$ (see (4.3)). In this case we write $\lambda_1 \sim \lambda_2$. We say that λ_1 *divides* λ_2 (or λ_2 is a *multiple* of λ_1) and write $\lambda_1 | \lambda_2$ if $\lambda_1 \sim \lambda_2$ and there exists $\lambda \in \Lambda$ such that $\lambda \sim \lambda_1$ and $\lambda_2 = \lambda\lambda_1$.

Let K be a difference-differential field ($Char\,K = 0$) with the basic set $\Delta \bigcup \sigma$ described above and let partition (3.1) of the set Δ be fixed. Let $K\{y_1, \dots, y_s\}$ be the ring of Δ-σ-polynomials over K and let ΛY denote the set of all elements λy_i $(\lambda \in \Lambda,\ 1 \leq i \leq s)$ called *terms*. Note that as a ring, $K\{y_1, \dots, y_s\} = K[\Lambda Y]$. Two terms $u = \lambda y_i$ and $v = \lambda' y_j$ are called *similar* if λ and λ' are similar; in this case we write $u \sim v$. If $u = \lambda y_i$ is a term and $\lambda' \in \Lambda$, we say that u is similar to λ' and write $u \sim \lambda'$ if $\lambda \sim \lambda'$. Furthermore, if $u, v \in \Lambda Y$, we say that u *divides* v or v *is a multiple of* u, if $u = \lambda y_i$, $v = \lambda' y_i$ for some y_i and $\lambda | \lambda'$.

Let us consider $p+1$ orders $<_1, \dots, <_p, <_\sigma$ on the set ΛY that correspond to the orders on the semigroup Λ (we use the same symbols for the orders on Λ and ΛY). These orders are defined as follows: $\lambda y_j <_i$ (or $<_\sigma$) $\lambda' y_k$ if and only if $\lambda <_i$ (respectively, $<_\sigma$)λ' in Λ or $\lambda = \lambda'$ and $j < k$ ($1 \leq i \leq p$, $1 \leq j, k \leq s$).

The order of a term $u = \lambda y_k$ and its orders with respect to the sets Δ_i ($1 \leq i \leq p$) and σ are defined as the corresponding orders of λ (we use the same notation $ord\, u$, $ord_i u$, and ord_σ for the corresponding orders).

If $A \in K\{y_1, \dots, y_s\} \setminus K$ and $1 \leq k \leq p$, then the highest with respect to $<_k$ term that appears in A is called the *k-leader* of the Δ-σ-polynomial A. It is denoted by $u_A^{(k)}$. The highest term of A with respect to $<_\sigma$ is called the *σ-leader* of A; it is denoted by v_A.

If A is written as a polynomial in v_A, $A = I_d(v_A)^d + I_{d-1}(v_A)^{d-1} + \cdots + I_0$, where all terms of $I_0, \dots, I_d$ are less than v_A with respect to $<_\sigma$, then I_d is said to be the *initial* of A. The partial derivative of A with respect to v_A, $\partial A / \partial v_A = dI_d(v_A)^{d-1} + (d-1)I_{d-1}(v_A)^{d-2} + \cdots + I_1$, is called the *separant* of A. The leading coefficient and the separant of a Δ-σ-polynomial A are denoted by I_A and S_A, respectively.

If $A, B \in K\{y_1, \dots, y_s\}$, then A is said to have lower rank than B (we write $rk\, A < rk\, B$) if either $A \in K$, $B \notin K$, or $(v_A, deg_{v_A} A, ord_1 u_A^{(1)}, \dots, ord_p u_A^{(p)})$ is less than $(v_B, deg_{v_B} B, ord_1 u_B^{(1)}, \dots, ord_p u_B^{(p)})$ with respect to the lexicographic order (v_A and v_B are compared with respect to $<_\sigma$).

If the vectors are equal (or $A, B \in K$) we say that A and B are of the same rank and write $rk\, A = rk\, B$.

Definition 5.1 *If $A, B \in K\{y_1, \dots, y_s\}$, then B is said to be reduced with respect to A if*

(i) B does not contain terms λv_A such that $\lambda \sim v_A$, $\lambda_\Delta \neq 1$, and $ord_i(\lambda u_A^{(i)}) \leq ord_i u_B^{(i)}$ for $i = 1, \dots, p$.

(ii) If B contains a term λv_A, where $\lambda \sim v_A$, $\lambda_\Delta = 1$, then either there exists j, $1 \leq j \leq p$, such that $ord_j u_B^{(j)} < ord_j(\lambda u_A^{(j)})$ or $ord_j(\lambda u_A^{(j)}) \leq ord_j u_B^{(j)}$ for all $j = 1, \dots, p$ and $deg_{\lambda v_A} B < deg_{v_A} A$.

If $B \in K\{y_1, \dots, y_s\}$, then B is said to be *reduced with respect to a set* $\Sigma \subseteq K\{y_1, \dots, y_s\}$ if B is reduced with respect to every element of Σ. A set $\Sigma \subseteq K\{y_1, \dots, y_s\}$ is called *autoreduced* if $\Sigma \bigcap K = \emptyset$ and every element of Σ is reduced with respect to any other element of this set.

The proof of the following lemma can be found in [5, Chapter 0, Section 17].

Lemma 5.2 *Let A be any infinite subset of the set $\mathbf{N}^m \times \mathbf{N}_n$ ($m, n \in \mathbf{N}$, $n \geq 1$). Then there exists an infinite sequence of elements of A, strictly increasing relative to the product order, in which every element has the same projection on $\mathbf{N}_n$.*

This lemma immediately implies the following statement that will be used below.

Lemma 5.3 *Let S be any infinite set of terms λy_j ($\lambda \in \Lambda, 1 \leq j \leq s$) in the ring $K\{y_1, \dots, y_s\}$. Then there exists an index j ($1 \leq j \leq s$) and an infinite sequence of terms $\lambda_1 y_j, \lambda_2 y_j, \dots, \lambda_k y_j, \dots$ such that $\lambda_k | \lambda_{k+1}$ for every $k = 1, 2, \dots$.*

Proposition 5.4 *Every autoreduced set is finite.*

PROOF. Suppose that Σ is an infinite autoreduced subset of $K\{y_1, \dots, y_s\}$. Then Σ must contain an infinite set $\Sigma' \subseteq \Sigma$ such that all Δ-σ-polynomials from Σ' have different σ-leaders similar to each other. Indeed, if it is not so, then there exists an infinite set $\Sigma_1 \subseteq \Sigma$ such that all Δ-σ-polynomials in Σ_1 have the same σ-leader v. By Lemma 5.2, the infinite set $\{(ord_1 u_A^{(1)}, \dots, ord_p u_A^{(p)}) | A \in \Sigma_1\}$ contains a nondecreasing infinite sequence $(ord_1 u_{A_1}^{(1)}, \dots, ord_p u_{A_1}^{(p)}) \leq_P (ord_1 u_{A_2}^{(1)}, \dots, ord_p u_{A_2}^{(p)}) \leq_P \dots$ ($A_1, A_2, \dots \in \Sigma_1$ and $\leq_P$ denotes the product order on $\mathbf{N}^p$) such that $v_{A_i} | v_{A_{i+1}}$. Since the sequence $\{deg_v A_i | i = 1, 2, \dots\}$ cannot be strictly decreasing, there exists two indices i and j such that $i < j$ and $deg_{v_{A_i}} A_i \leq deg_{v_{A_j}} A_j$. We obtain that A_j is reduced with respect to A_i that contradicts the fact that Σ is an autoreduced set.

Thus, we can assume that all Δ-σ-polynomials of our infinite autoreduced set Σ have distinct σ-leaders. By Lemma 5.3, there exists an infinite sequence $B_1, B_2, \dots$ of elements of Σ such that $v_{B_i} | v_{B_{i+1}}$ for all $i = 1, 2, \dots$. Let $k_{ij} = ord_\sigma v_{B_i}$ and $l_{ij} = ord_j u_{B_i}^{(j)}$ ($1 \leq j \leq p$). Obviously, $l_{ij} \geq k_{ij}$ ($i = 1, 2, \dots; j = 1, \dots, p$), so that $\{(l_{i1} - k_{i1}, \dots, l_{ip} - k_{ip}) | i = 1, 2, \dots\} \subseteq \mathbf{N}^p$. By Lemma 5.2, there exists an infinite sequence of indices $i_1 < i_2 < \dots$ such that $(l_{i_1 1} - k_{i_1 1}, \dots, l_{i_1 p} - k_{i_1 p}) \leq_P (l_{i_2 1} - k_{i_2 1}, \dots, l_{i_2 p} - k_{i_2 p}) \leq_P \dots$. Then for any $j = 1, \dots, p$, we have $ord_j\, (\frac{v_{B_{i_2}}}{v_{B_{i_1}}} u_{B_{i_1}}^{(j)}) = k_{i_2 j} - k_{i_1 j} + l_{i_1 j} \leq k_{i_2 j} + l_{i_2 j} - k_{i_2 j} = l_{i_2 j} = ord_j u_{B_{i_2}}^{(j)}$, so that B_{i_2} contains a term $\lambda v_{B_{i_1}} = v_{B_{i_2}}$ such that $\lambda \neq 1$ and $ord_j(\lambda u_{B_{i_1}}^{(j)}) \leq ord_j u_{B_{i_2}}^{(j)}$ for $j = 1, \dots, p$. Thus, the Δ-σ-polynomial B_{i_2} is reduced with respect to B_{i_1} that contradicts the fact that Σ is an autoreduced set.

Throughout the rest of the paper, while considering autoreduced sets in the ring $K\{y_1, \dots, y_s\}$ we always assume that their elements are arranged in order of increasing rank. (Therefore, if we consider an autoreduced set of Δ-polynomials $\Sigma = \{A_1, \dots, A_r\}$, then $rk\, A_1 < \dots < rk\, A_r$).

Theorem 5.5 *Let $\Sigma = \{A_1, \dots, A_d\}$ be an autoreduced set in the ring $R = K\{y_1, \dots, y_s\}$ and let I_k and S_k denote the initial and separant of A_k, respectively. Furthermore, let $I(\Sigma) = \{X \in K\{y_1, \dots, y_s\} \,|\, X = 1$ or X is a product of finitely many elements of the form $\gamma(I_k)$ and $\gamma'(S_k)$ where $\gamma, \gamma' \in \Lambda_\sigma\}$. Then for any Δ-σ-polynomial B, there exist $B_0 \in K\{y_1, \dots, y_s\}$ and $J \in I(\Sigma)$ such that B_0 is reduced with respect to Σ and $JB \equiv B_0\, (mod [\Sigma])$ (that is, $JB - B_0 \in [\Sigma]$).*

PROOF. If B is reduced with respect to Σ, the statement is obvious (one can set $B_0 = B$). Suppose that B is not reduced with respect to Σ. Let $u_i^{(j)}$ and v_i $(1 \leq j \leq p,\ 1 \leq i \leq d)$ be the leaders of the element A_i relative to the orders $<_j$ and $<_\sigma$, respectively. In what follows, a term w_H, that appears in a Δ-σ-polynomial $H \in R$, will be called a Σ-leader of H if w_H is the greatest (with respect to $<_\sigma$) term among all terms λv_i $(1 \leq i \leq d)$ such that $\lambda \sim v_i$, λv_i appears in H and either $\lambda_\Delta \neq 1$ and $ord_j(\lambda u_i^{(j)}) \leq ord_j u_H^{(j)}$ for $j = 1, \dots, p$, or $\lambda_\Delta = 1$, $ord_j(\lambda u_i^{(j)}) \leq ord_j u_H^{(j)}$ $(1 \leq j \leq p)$, and $deg_{v_i} A_i \leq deg_{\lambda v_i} H$.

Let w_B be the Σ-leader of B. Then the Δ-σ-polynomial B can be written as $B = B' w_B^r + B''$ where B' does not contain w_B and $deg_{w_B} B'' < r$. Let $w_B = \lambda v_i$ for some i $(1 \leq i \leq d)$ and for some $\lambda \in \Lambda$, $\lambda \sim w_B \sim v_i$, such that $ord_j(\lambda u_i^{(j)}) \leq ord_j u_B^{(j)}$ for $j = 1, \dots, p$. Without loss of generality we may assume that i corresponds to the maximum (with respect to the order $<_\sigma$) σ-leader v_i in the set of all σ-leaders of elements of Σ.

Suppose, first, that $\lambda_\Delta \neq 1$ (and $ord_j(\lambda u_i^{(j)}) \leq ord_j u_B^{(j)}$ for $j = 1, \dots, p$). Then $\lambda_\Delta A_i - S_i \lambda_\Delta v_i$ has lower rank than $\lambda_\Delta v_i$, hence $T = \lambda A_i - \lambda_\sigma(S_i)\lambda v_i$ has lower rank than $\lambda v_i = w_B$. Also, $(\lambda_\sigma(S_i))^r B = (\lambda_\sigma(S_i)\lambda v_i)^r B' + (\lambda_\sigma(S_i))^r B'' = (\lambda A_i - T)^r B' + (\lambda_\sigma(S_i))^r B''$. Setting $B^{(1)} = B'(-T)^r + (\lambda_\sigma(S_i))^r B''$ we obtain that $B^{(1)} \equiv B\, mod[\Sigma]$, $B^{(1)}$ is reduced with respect to Σ in the sense of Definition 5.1, $B^{(1)}$ does not contain any Σ-leader, which is greater than w_B with respect to $<_\sigma$, and $deg_{w_B} B^{(1)} < r$.

Now let $\lambda_\Delta = 1$, $ord_j(\lambda u_i^{(j)}) \leq ord_j u_B^{(j)}$ $(1 \leq j \leq p)$, and $r_i < r$ where $r_i = deg_{v_i} A_i$. Then the Δ-σ-polynomial $(\lambda I_i)B - w_B^{r - r_i}(\lambda A_i)B'$ has all the properties of $B^{(1)}$ mentioned above. Repeating the described procedure, we arrive at a desired Δ-σ-polynomial B_0 that does not contain any Σ-leader (so it is reduced with respect to Σ) and satisfies the condition $JB \equiv B_0\, (mod[\Sigma])$ where $J = 1$ or J is a product of finitely many elements of the form $\gamma(I_k)$ and $\gamma'(S_k)$ $(\gamma, \gamma' \in \Lambda_\sigma)$.

With the notation of the last theorem, we say that the Δ-σ-polynomial B *reduces to* B_0 modulo Σ.

Definition 5.6 *Let* $\Sigma = \{A_1, \dots, A_d\}$ *and* $\Sigma' = \{B_1, \dots, B_e\}$ *be two autoreduced sets in the ring of differential polynomials* $K\{y_1, \dots, y_s\}$*. An autoreduced set* Σ *is said to have lower rank than* Σ' *if one of the following two cases holds:*

(1) There exists $k \in \mathbf{N}$ *such that* $k \leq \min\{d, e\}$, $rk\, A_i = rk\, B_i$ *for* $i = 1, \dots, k-1$ *and* $rk\, A_k < rk\, B_k$*.*

(2) $d > e$ *and* $rk\, A_i = rk\, B_i$ *for* $i = 1, \dots, e$*.*

If $d = e$ *and* $rk\, A_i = rk\, B_i$ *for* $i = 1, \dots, d$*, then* Σ *is said to have the same rank as* Σ'*.*

Proposition 5.7 *In every nonempty family of autoreduced sets of differential polynomials there exists an autoreduced set of lowest rank.*

PROOF. Let Φ be any nonempty family of autoreduced sets in the ring $K\{y_1, \dots, y_s\}$. Let us inductively define an infinite descending chain of subsets of Φ as follows: $\Phi_0 = \Phi$, $\Phi_1 = \{\Sigma \in \Phi_0 | \Sigma$ contains at least one element and the first element of Σ is of lowest possible rank$\}$, ..., $\Phi_k = \{\Sigma \in \Phi_{k-1} | \Sigma$ contains at least k elements and the kth element of Σ is of lowest possible rank$\}$, ... It is clear that if A and B are any two Δ-σ-polynomials in the same set Φ_k, then $v_A = v_B$, $deg_{v_A} A = deg_{v_B} B$, and $ord_i u_A^{(i)} = ord_i u_B^{(i)}$ for $i = 1, \dots, p$. Therefore, if all sets Φ_k are nonempty, then the set $\{A_k | A_k$ is the kth element of some autoreduced set in $\Phi_k\}$ would be an infinite autoreduced set, and this would contradict Lemma 5.3. Thus, there is the smallest positive integer k such that Φ_k is empty. It is clear that every element of Φ_{k-1} is an autoreduced set of lowest rank in Φ.

Let J be any ideal of the ring $K\{y_1, \dots, y_s\}$. Since the set of all autoreduced subsets of J is not empty (if $A \in J$, then $\{A\}$ is an autoreduced subset of J), the last statement shows that the ideal J contains an autoreduced subset of lowest rank. Such an autoreduced set is called a *characteristic set* of the ideal J.

Proposition 5.8 *Let $\Sigma = \{A_1, \dots, A_d\}$ be a characteristic set of a Δ-σ-ideal J of the ring $R = K\{y_1, \dots, y_s\}$. Then an element $B \in R$ is reduced with respect to the set Σ if and only if $B = 0$.*

PROOF. First of all, note that if $B \neq 0$ and $rk\, B < rk\, A_1$, then $rk\, \{B\} < rk\, \Sigma$ that contradicts the fact that Σ is a characteristic set of the ideal J. Let $rk\, B > rk\, A_1$ and let $A_1, \dots, A_j$ $(1 \leq j \leq d)$ be all elements of Σ whose rank is lower that the rank of B. Then the set $\Sigma' = \{A_1, \dots, A_j, B\}$ is autoreduced. Indeed, by the conditions of the theorem, Δ-σ-polynomials $A_1, \dots, A_j$ are reduced with respect to each other and B is reduced with respect to the set $\{A_1, \dots, A_j\}$. Furthermore, each A_i $(1 \leq i \leq j)$ is reduced with respect to B because $rk\, A_i < rk\, B$. Since $rk\, \Sigma' < rk\, \Sigma$, Σ is not a characteristic set of J that contradicts the conditions of the theorem. Thus, $B = 0$.

Now we can introduce the concept of a coherent autoreduced set of a linear Δ-σ-ideal of $K\{y_1, \dots, y_s\}$ (that is, a Δ-σ-ideal generated by a finite set of linear Δ-σ-polynomials) in the same way as it is defined in the case of difference polynomials (see [6, Section 6.5]): an autoreduced set $\Sigma = \{A_1, \dots, A_d\} \subseteq K\{y_1, \dots, y_s\}$ consisting of linear Δ-σ-polynomials is called *coherent* if it satisfies the following two conditions:

(i) λA_i reduces to zero modulo Σ for any $\lambda \in \Lambda$, $1 \leq i \leq d$.

(ii) If $v_{A_i} \sim v_{A_j}$ and $w = \lambda v_{A_i} = \lambda' v_{A_j}$, where $\lambda \sim \lambda' \sim v_{A_i} \sim v_{A_j}$, then the Δ-σ-polynomial $(\lambda' I_{A_j})(\lambda A_i) - (\lambda I_{A_i})(\lambda' A_j)$ reduces to zero modulo Σ.

The following two propositions can be proved precisely in the same way as the corresponding statements for difference polynomials, see [6, Theorem 6.5.3 and Corollary 6.5.4]).

Proposition 5.9 *Any characteristic set of a linear Δ-σ-ideal of the ring of Δ-σ-polynomials $K\{y_1, \dots, y_s\}$ is a coherent autoreduced set. Conversely, if Σ is a* *coherent autoreduced set in $K\{y_1, \dots, y_s\}$ consisting of linear Δ-σ-polynomials, then Σ is a characteristic set of the linear Δ-σ-ideal $[\Sigma]$.*

Proposition 5.10 *Let us consider a partial order $\preccurlyeq$ on $K\{y_1, \dots, y_s\}$ such that $A \preccurlyeq B$ if and only if $v_A | v_B$. Let A be a linear Δ-σ-polynomial in $K\{y_1, \dots, y_s\} \setminus K$. Then the set of all minimal with respect to $\preccurlyeq$ elements of the set $\{\lambda A \,|\, \lambda \in \Lambda\}$ is a characteristic set of the Δ-σ-ideal $[A]$.*

Now we are ready to prove Theorem 3.1.

PROOF. Let $L = K\langle \eta_1, \dots, \eta_s \rangle$ be a Δ-σ-field extension of K generated by a finite set $\eta = \{\eta_1, \dots, \eta_s\}$. Then there exists a natural Δ-σ-homomorphism Υ_η of the ring of Δ-σ-polynomials $K\{y_1, \dots, y_s\}$ onto the Δ-σ-subring $K\{\eta_1, \dots, \eta_s\}$ of L such that $\Upsilon_\eta(a) = a$ for any $a \in K$ and $\Upsilon_\eta(y_j) = \eta_j$ for $j = 1, \dots, n$. (If $A \in K\{y_1, \dots, y_s\}$, then $\Upsilon_\eta(A)$ is called the *value* of A at η and it is denoted by $A(\eta)$.) Obviously, the kernel P of the Δ-σ-homomorphism Υ_η is a prime Δ-σ-ideal of the ring $K\{y_1, \dots, y_s\}$. This ideal is called the *defining* ideal of η over K or the defining ideal of the Δ-σ-field extension $G = K\langle \eta_1, \dots, \eta_s \rangle$. It is easy to see that if the quotient field Q of the factor ring $\bar{R} = K\{y_1, \dots, y_s\}/P$ is considered as a Δ-σ-field (where $\delta(\frac{f}{g}) = \frac{f\delta(g) - f\delta(g)}{g^2}$ and $\tau(\frac{f}{g}) = \frac{\tau(f)}{\tau(g)}$ for any $f, g \in \bar{R}$, $\delta \in \Delta$, $\tau \in \sigma^*$), then this quotient field is naturally Δ-σ-isomorphic to the field L. The Δ-σ-isomorphism of Q onto L is identity on K and maps the images of the Δ-σ-indeterminates $y_1, \dots, y_s$ in the factor ring $\bar{R}$ onto the elements $\eta_1, \dots, \eta_s$, respectively.

Let $\Sigma = \{A_1, \dots, A_d\}$ be a characteristic set of the defining Δ-σ-ideal P. For any $r_1, \dots, r_{p+1} \in \mathbf{N}$, let us set $U_{r_1 \dots r_{p+1}} = \{u \in \Lambda Y | ord_i u \leq r_i$ for $i = 1, \dots, p{+}1$ and either u is not a multiple of any v_{A_i} or for every $\lambda \in \Lambda$, $A \in \Sigma$ such that $u = \lambda v_A$ and $\lambda \sim v_A$, there exists $j \in \{1, \dots, p\}$ such that $ord_j(\lambda u_A^{(j)}) > r_j\}$. (For shortness, here and below we sometimes write $\leq_{i+1}$ instead of $\leq_\sigma$.)

We are going to show that the set $\bar{U}_{r_1 \dots r_p} = \{u(\eta) | u \in U_{r_1 \dots r_p}\}$ is a transcendence basis of the field $K(\bigcup\limits_{j=1}^{n} \Lambda(r_1, \dots, r_{p+1})\eta_j)$ over K.

First of all, let us show that the set $\bar{U}_{r_1 \dots r_{p+1}}$ is algebraically independent over K. Let g be a polynomial in k variables ($k \in \mathbf{N}, k \geq 1$) such that $g(u_1(\eta), \dots, u_k(\eta)) = 0$ for some elements $u_1, \dots, u_k \in U_{r_1 \dots r_{p+1}}$. Then the Δ-σ-polynomial $\bar{g} = g(u_1, \dots, u_k)$ is reduced with respect to Σ. (Indeed, if g contains a term $u = \lambda v_{A_i}$ with $\lambda \in \Lambda$, $\lambda \sim v_{A_i}$ $(1 \leq i \leq d)$, then there exists $k \in \{1, \dots, p\}$ such that $ord_k(\lambda u_{A_i}^{(k)}) > r_k \geq ord_k u_{\bar{g}}^{(k)})$. Since $\bar{g} \in P$, Proposition 5.8 implies that $\bar{g} = 0$. Thus, the set $\bar{U}_{r_1 \dots r_p}$ is algebraically independent over K.

Now, let us prove that every element $\lambda\eta_j$ $(1 \leq j \leq s, \lambda \in \Lambda(r_1, \dots, r_{p+1}))$ is algebraic over the field $K(\bar{U}_{r_1, \dots, r_{p+1}})$. Let $\lambda\eta_j \notin \bar{U}_{r_1, \dots, r_{p+1}}$ (if $\lambda\eta_j \in \bar{U}_{r_1, \dots, r_{p+1}}$, the statement is obvious). Then $\lambda y_j \notin U_{r_1, \dots, r_p}$ whence λy_j is equal to some term of the form $\lambda' v_{A_i}$ where $\lambda' \in \Lambda$, $\lambda \sim v_{A_i}$ $(1 \leq i \leq d)$, and $ord_k(\lambda' u_{A_i}^{(k)}) \leq r_k$ for $k = 1, \dots, p$. Let us represent A_i as a polynomial in v_{A_i}: $A_i = I_0 {(v_{A_i})}^e + I_1 {(v_{A_i})}^{e-1} + \dots + I_e$, where $I_0, I_1, \dots I_e$ do not contain v_{A_i} (therefore, all terms in these Δ-σ-polynomials are lower than v_{A_i} with respect to the order $<_\sigma$). Since $A_i \in P$,

$$A_i(\eta) = I_0(\eta)(v_{A_i}(\eta))^e + I_1(\eta)(v_{A_i}(\eta))^{e-1} + \cdots + I_e(\eta) = 0 \qquad (5.\ 1)$$

It is easy to see that the Δ-σ-polynomials I_0 and $S_{A_i} = \partial A_i/\partial v_{A_i}$ are reduced with respect to any element of the set Σ. Applying Proposition 5.8 we obtain that $I_0 \notin P$ and $S_{A_i} \notin P$ whence $I_0(\eta) \neq 0$ and $S_{A_i}(\eta) \neq 0$. Now, if we apply λ' to both sides of equation (5.1), the resulting equation will show that the element $\lambda' v_{A_i}(\eta) = \lambda\eta_j$ is algebraic over the field $K(\{\bar{\lambda}\eta_l | ord_i\bar{\lambda} \leq r_i$ for $i = 1, \dots, p+1; 1 \leq l \leq s$, and $\bar{\lambda}y_l <_\sigma \lambda' u^{(1)}_{A_i} = \lambda y_j\})$. Now, the induction on the set of terms ΛY ordered by the relation $<_\sigma$ completes the proof of the fact that $\bar{U}_{r_1 \dots r_{p+1}}(\eta)$ is a transcendence basis of the field $K(\bigcup_{j=1}^{s} \Lambda(r_1, \dots, r_{p+1})\eta_j)$ over K.

Let $U^{(1)}_{r_1 \dots r_{p+1}} = \{u \in \Lambda Y | ord_i u \leq r_i$ for $i = 1, \dots, p{+}1$ and u is not a multiple of any v_{A_j}, $j = 1, \dots, d\}$ and $U^{(2)}_{r_1 \dots r_{p+1}} = \{u \in \Lambda Y | ord_i u \leq r_i$ for $i = 1, \dots, p{+}1$ and there exists at least one pair i, j $(1 \leq i \leq p,\ 1 \leq j \leq d)$ such that $u = \lambda v_{A_j}$, $\lambda \sim v_{A_j}$, and $ord_i(\lambda u^{(i)}_{A_j}) > r_i\}$. Clearly, $U_{r_1 \dots r_{p+1}} = U^{(1)}_{r_1 \dots r_{p+1}} \bigcup U^{(2)}_{r_1 \dots r_{p+1}}$ and $U^{(1)}_{r_1 \dots r_{p+1}} \bigcap U^{(2)}_{r_1 \dots r_{p+1}} = \emptyset$.

By Theorem 4.6, there exists a numerical polynomial $\phi(t_1, \dots, t_{p+1})$ in $p{+}1$ variables $t_1, \dots, t_{p+1}$ such that $\phi(r_1, \dots, r_{p+1}) = Card\, U^{(1)}_{r_1 \dots r_{p+1}}$ for all sufficiently large $(r_1 \dots r_{p+1}) \in \mathbf{N}^{p+1}$, $deg_{t_i}\phi \leq m_i$ for $i = 1, \dots, p$, and $deg_{t_{p+1}}\phi \leq n$. Thus, in order to complete the proof of the theorem, we need to show that there exists a numerical polynomial $\psi(t_1, \dots, t_{p+1})$ in $p{+}1$ variables $t_1, \dots, t_{p+1}$ such that $\psi(r_1, \dots, r_{p+1}) = Card\, U^{(2)}_{r_1 \dots r_{p+1}}$ for all sufficiently large $(r_1 \dots r_{p+1}) \in \mathbf{N}^{p+1}$, $deg_{t_i}\psi \leq m_i$ for $i = 1, \dots, p$, and $deg_{t_{p+1}}\psi \leq n$.

Let $ord_i v_{A_j} = a_{ij}$, $ord_i u^{(i)}_{A_j} = b_{ij}$, and $ord_\sigma v_{A_j} = c_j$ for $i = 1, \dots, p; j = 1, \dots, d$ (clearly, $a_{ij} \leq b_{ij}$ for $i = 1, \dots, p$; $j = 1, \dots, d$). Furthermore, for any $q = 1, \dots, p$ and for any integers $k_1, \dots, k_q$ such that $1 \leq k_1 < \cdots < k_q \leq p$, let $V_{j;k_1,\dots,k_q}(r_1, \dots, r_{p+1}) = \{\lambda v_{A_j} | \lambda \sim v_{A_j}$, $ord_i\lambda \leq r_i - a_{ij}$ for $i = 1, \dots, p$, $ord_\sigma\lambda \leq r_{p+1} - c_j$, and $ord_k\lambda > r_k - b_{kj}$ if and only if k is equal to one of the numbers $k_1, \dots, k_q\}$. Using Theorem 4.3(iii) we obtain that $Card\, V_{j;k_1,\dots,k_q}(r_1, \dots, r_{p+1}) = \phi_{j;k_1,\dots,k_q}(r_1, \dots, r_{p+1})$, where $\phi_{j;k_1,\dots,k_q}(t_1, \dots, t_{p+1})$ is a numerical polynomial in p variables $t_1, \dots, t_{p+1}$ defined by the formula

$$\phi_{j;k_1,\dots,k_q}(t_1, \dots, t_p) = \binom{t_1 + m_1 - a_{1j}}{m_1} \cdots \binom{t_{k_1-1} + m_{k_1-1} - a_{k_1-1,j}}{m_{k_1-1}}$$

$$\left[\binom{t_{k_1} + m_{k_1} - a_{k_1,j}}{m_{k_1}} - \binom{t_{k_1} + m_{k_1} - b_{k_1,j}}{m_{k_1}}\right] \binom{t_{k_1+1} + m_{k_1+1} - a_{k_1+1,j}}{m_{k_1+1}}$$

$$\cdots \binom{t_{k_q-1} + m_{k_q-1} - a_{k_q-1,j}}{m_{k_q-1}} \left[\binom{t_{k_q} + m_{k_q} - a_{k_q,j}}{m_{k_q}} - \binom{t_{k_q} + m_{k_q} - b_{k_q,j}}{m_{k_q}}\right]$$

$$\dots\binom{t_p+m_p-a_{pj}}{m_p}\binom{t_{p+1}+n-c_j}{n} \qquad (5.\ 2)$$

Clearly, $deg_{t_i}\phi_{j;k_1,\dots,k_q} \leq m_i$ for $i=1,\dots,p$ and $deg_{t_{p+1}}\phi_{j;k_1,\dots,k_q} \leq n$.

Now, for any $j=1,\dots,d$, let $V_j(r_1,\dots,r_{p+1}) = \{\lambda v_{A_j} | \lambda \sim v_{A_j},\ ord_i\lambda \leq r_i - a_{ij}$ for $i=1,\dots,p$, $ord_\sigma\lambda \leq r_{p+1}-c_j$, and there exists $k \in \mathbf{N}$, $1 \leq k \leq p$, such that $ord_k\lambda > r_k - b_{kj}\}$. Then the combinatorial principle of inclusion and exclusion implies that $Card\, V_j(r_1,\dots,r_{p+1}) = \phi_j(r_1,\dots,r_{p+1})$, where $\phi_j(t_1,\dots,t_{p+1})$ is a numerical polynomial in $p+1$ variables $t_1,\dots,t_{p+1}$ defined by the formula

$$\phi_j(t_1,\dots,t_{p+1}) = \sum_{k_1=1}^{p}\phi_{j;k_1}(t_1,\dots,t_{p+1}) - \sum_{1\leq k_1<k_2\leq p}\phi_{j;k_1,k_2}(t_1,\dots,t_{p+1}) + \dots +$$

$$(-1)^{\nu-1}\sum_{1\leq k_1<\dots k_\nu\leq p}\phi_{j;k_1,\dots,k_\nu}(t_1,\dots,t_{p+1}) + \dots + (-1)^p\phi_{j;1,2,\dots,p}(t_1,\dots,t_{p+1}). \qquad (5.\ 3)$$

It is easy to see that $deg_{t_i}\phi_j \leq m_i$ for $i=1,\dots,p$ and $deg_{t_{p+1}}\phi_j \leq n$.

Applying the principle of inclusion and exclusion we obtain that

$$Card\, U^{(2)}_{r_1\dots r_{p+1}} = Card\bigcup_{j=1}^{d}V_j(r_1,\dots,r_{p+1}) = \sum_{j=1}^{d}Card\, V_j(r_1,\dots,r_{p+1})$$

$$-\sum_{1\leq j_1<j_2\leq d}Card\,(V_{j_1}(r_1,\dots,r_{p+1})\bigcap V_{j_2}(r_1,\dots,r_{p+1})) + \dots +$$

$$(-1)^{d-1}Card\bigcap_{\nu=1}^{d}V_\nu(r_1,\dots,r_{p+1}), \qquad (5.\ 4)$$

so it is sufficient to prove that for any $s=1,\dots,d$ and for any indices $j_1,\dots,j_s$, $1 \leq j_1 < \dots < j_s \leq d$, $Card\,(V_{j_1}(r_1,\dots,r_{p+1})\bigcap\dots\bigcap V_{j_s}(r_1,\dots,r_{p+1})) = \phi_{j_1,\dots,j_s}(r_1,\dots,r_{p+1})$, where $\phi_{j_1,\dots,j_s}(t_1,\dots,t_{p+1})$ is a numerical polynomial in $p+1$ variables $t_1,\dots,t_{p+1}$ such that $deg_{t_i}\phi_{j_1,\dots,j_s} \leq m_i$ for $i=1,\dots,p$ and $deg_{t_{p+1}}\phi_{j_1,\dots,j_s} \leq n$.

It is clear that the intersection $V_{j_1}(r_1,\dots,r_{p+1})\bigcap\dots\bigcap V_{j_s}(r_1,\dots,r_{p+1})$ is not empty (therefore, $\phi_{j_1,\dots,j_s} \neq 0$) if and only if the leaders $v_{A_{j_1}},\dots,v_{A_{j_s}}$ contain the same Δ-σ-indeterminate y_i $(1 \leq i \leq s)$ and they are all similar to each other. Let us consider such an intersection and let $v(j_1,\dots,j_s) = lcm(v_{A_{j_1}},\dots,v_{A_{j_s}})$ (this least common multiple of similar to each other terms is defined in a natural way, as the smallest with respect to $<_\sigma$ common multiple of $v_{A_{j_1}},\dots,v_{A_{j_s}}$). Let elements $\gamma_1,\dots,\gamma_s \in \Lambda$ be defined by the conditions $v(j_1,\dots,j_s) = \gamma_\nu v_{A_{j_\nu}}$ and $\gamma_\nu \sim v_{A_{j_\nu}}$ $(\nu=1,\dots,s)$.

Then $V_{j_1}(r_1,\dots,r_{p+1})\bigcap\dots\bigcap V_{j_s}(r_1,\dots,r_{p+1})$ is the set of all terms $u = \lambda v(j_1,\dots,j_s)$ such that $u \sim v(j_1,\dots,j_s)$, $ord_i u \leq r_i$ (that is, $ord_i\lambda \leq r_i -$

$ord_i v(j_1, \dots, j_s))$ for $i = 1, \dots, p$, $ord_\sigma u \leq r_{p+1}$, and for any $l = 1, \dots, s$, there exists at least one index $k \in \{1, \dots, p\}$ such that $ord_k(\lambda\gamma_l u^{(k)}_{A_{j_l}}) > r_k$ (i. e., $ord_k\lambda > r_k - ord_k v(j_1, \dots, j_s) - ord_k u^{(k)}_{A_{j_l}} + ord_k v_{A_{j_l}}$).

Setting $c^{(i)}_{j_1,\dots,j_s} = ord_i v(j_1, \dots, j_s)$ $(1 \leq i \leq p)$, $c^{(p+1)}_{j_1,\dots,j_s} = ord_\sigma v(j_1, \dots, j_s)$ and applying the principle of inclusion and exclusion once again, we obtain that $Card \bigcap_{\nu=1}^{s} V_{j_\nu}(r_1, \dots, r_{p+1})$ is an alternating sum of terms of the form $Card\, W(j_1, \dots, j_s; k_{11}, k_{12}, \dots, k_{1q_1}, k_{21}, \dots, k_{sq_s}; r_1, \dots, r_{p+1})$ where $W(j_1, \dots, j_s; k_{11}, k_{12}, \dots, k_{1q_1}, k_{21}, \dots, k_{sq_s}; r_1, \dots, r_{p+1})$ denotes the set

$\{\lambda \in \Lambda | \lambda \sim v(j_1, \dots, j_s), ord_i\lambda \leq r_i - c^{(i)}_{j_1,\dots,j_s}$ for $i = 1, \dots, p$, $ord_\sigma\lambda \leq r_{p+1} - c^{(p+1)}_{j_1,\dots,j_s}$ and for any $l = 1, \dots, s, 1 \leq k \leq p$, $ord_k\lambda > r_k - c^{(k)}_{j_1,\dots,j_s} + a_{kj_l} - b_{kj_l}$ if and only if k is equal to one of the numbers $k_{l1}, \dots k_{lq_l}\}$

$(q_1, \dots, q_s$ are some positive integers in the set $\{1, \dots, p\}$ and $\{k_{i\nu} | 1 \leq i \leq s, 1 \leq \nu \leq q_s\}$ is a family of integers such that $1 \leq k_{i1} < k_{i2} < \dots < k_{iq_i} \leq p$ for $i = 1, \dots, s)$.

Thus, it remains to show that $Card\, W(j_1, \dots, j_s; k_{11}, \dots, k_{sq_s}; r_1, \dots, r_{p+1}) = \psi^{j_1,\dots,j_s}_{k_{11},\dots,k_{sq_s}}(r_1, \dots, r_{p+1})$ where $\psi^{j_1,\dots,j_s}_{k_{11},\dots,k_{sq_s}}(t_1, \dots, t_{p+1})$ is a numerical polynomial in $p+1$ variables $t_1, \dots, t_{p+1}$ whose degrees with respect to t_i $(1 \leq i \leq p)$ and t_{p+1} do not exceed m_i and n, respectively. But this is almost evident: as in the evaluation of $Card\, V_{j;k_1,\dots,k_q}(r_1, \dots, r_{p+1})$ (when we use Theorem 4.3 (iii) to get formula (5.3)), we see that $Card\, W(j_1, \dots, j_s; k_{11}, \dots, k_{sq_s}; r_1, \dots, r_p)$ is a product of terms of the form $\binom{r_{p+1} + n - c^{(p+1)}_{j_1,\dots,j_s}}{n}$ or $\binom{r_k + m_k - c^{(k)}_{j_1,\dots,j_s} - S_k}{m_k}$ (the last term corresponds to an integer k such that $1 \leq k \leq p$ and $k \neq k_{i\nu}$ for any $i = 1, \dots, s, \nu = 1, \dots, q_s$; the number S_k is $\max\{b_{kj_l} - a_{kj_l} | 1 \leq l \leq s\}$) or $\left[\binom{r_k + m_k - c^{(k)}_{j_1,\dots,j_s}}{m_k} - \binom{r_k + m_k - c^{(k)}_{j_1,\dots,j_s} - T_k}{m_k}\right]$ (such a term appears in the product if k is equal to some $k_{i\nu}$ $(1 \leq i \leq s, 1 \leq \nu \leq q_s)$. In this case, if $k_{i_1\nu_1}, \dots, k_{i_e\nu_e}$ are all elements of the set $\{k_{i\nu} | 1 \leq i \leq s, 1 \leq \nu \leq q_s\}$ that are equal to k $(1 \leq e \leq s, 1 \leq i_1 < \dots < i_e \leq s)$, then T_k is defined as $\min\{b_{kj_{i_\lambda}} - a_{kj_{i_\lambda}} | 1 \leq \lambda \leq l\}$).

The corresponding numerical polynomial $\psi^{j_1,\dots,j_s}_{k_{11},\dots,k_{sq_s}}(t_1, \dots, t_{p+1})$ is a product of p "elementary" numerical polynomials $f_1, \dots, f_p$ where f_k $(1 \leq k \leq p)$ is a polynomial of the form $\binom{t_{p+1} + n - c^{(p+1)}_{j_1,\dots,j_s}}{n}$ or $\binom{t_k + m_k - c^{(k)}_{j_1,\dots,j_s} - S_k}{m_k}$ or

$\left[\binom{t_k + m_k - c^{(k)}_{j_1,\dots,j_s}}{m_k} - \binom{t_k + m_k - c^{(k)}_{j_1,\dots,j_s} - T_k}{m_k}\right]$ $(1 \leq k \leq p)$. Since the degree of such a product with respect to any variable t_i $(1 \leq i \leq p)$ and t_{p+1} does not exceed m_i and n, respectively, this completes the proof of the first two parts of Theorem 3.1.

In order to prove the last part of the theorem, suppose that $\zeta = \{\zeta_1, \dots, \zeta_q\}$

is another system of Δ-σ-generators of L/K, that is, $L = K\langle\eta_1, \dots, \eta_s\rangle = K\langle\zeta_1, \dots, \zeta_q\rangle$). Let

$$\Phi_\zeta(t_1, \dots, t_{p+1}) = \sum_{i_1=0}^{m_1} \dots \sum_{i_p=0}^{m_p} b_{i_1 \dots i_{p+1}} \binom{t_1 + i_1}{i_1} \dots \binom{t_{p+1} + i_{p+1}}{i_{p+1}}$$

be the dimension polynomial of our Δ-σ-field extension associated with the system of generators ζ. Then there exist $h_1, \dots, h_{p+1} \in \mathbf{N}$ such that $\eta_i \in K(\bigcup_{j=1}^{q} \Lambda(h_1, \dots, h_{p+1})\zeta_j)$ and $\zeta_k \in K(\bigcup_{j=1}^{s} \Lambda(h_1, \dots, h_{p+1})\eta_j)$ for any $i = 1, \dots, s$ and $k = 1, \dots, q$, whence $\Phi_\eta(r_1, \dots, r_{p+1}) \leq \Phi_\zeta(r_1+h_1, \dots, r_{p+1}+h_{p+1})$ and $\Phi_\zeta(r_1, \dots, r_{p+1}) \leq \Phi_\eta(r_1 + h_1, \dots,$
$r_{p+1} + h_{p+1})$ for all sufficiently large $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$. Now the statement of the third part of Theorem 3.1 follows from the fact that for any element $(k_1, \dots, k_{p+1}) \in E'_\eta$, the term $\binom{t_1 + k_1}{k_1} \dots \binom{t_{p+1} + k_{p+1}}{k_{p+1}}$ appears in $\Phi_\eta(t_1, \dots, t_{p+1})$ and $\Phi_\zeta(t_1, \dots, t_{p+1})$ with the same coefficient $a_{k_1 \dots k_{p+1}}$.

Let us consider some examples of computation of Δ-σ-dimension polynomials.

Example 5.11 Consider a Δ-σ-field extension $L = K\langle\eta\rangle$ where $\Delta = \{\delta\}$, $\sigma = \{\alpha\}$, and the Δ-σ-generator of L/K satisfies the defining equation

$$\delta\eta - \alpha\eta - a = 0 \tag{5. 5}$$

where $a \in K$.

Since the equation is linear, the corresponding defining Δ-σ-ideal of the ring of Δ-σ-polynomials $K\{y\}$ is $P = [\alpha y - \delta y - a]$ (the fact that any linear ideal is prime is well-known, see, for example, [5, Chapter IV, Section 5]). By Proposition 5.10, the Δ-σ-polynomials $A = \alpha y - \delta y - a$ and $-\alpha^{-1}A = \alpha^{-1}\delta y - y - \alpha^{-1}(a)$ form a characteristic set of the ideal P. Using the notation of the proof of Theorem 3.1 and applying the procedure described in this proof together with Theorem 4.3(iii), and formula (4.4), we obtain that $Card\, U^{(1)}_{r_1 r_2} = r_1 + r_2 + 1$ and $Card\, U^{(2)}_{r_1 r_2} = r_2$ for all sufficiently large $(r_1, r_2) \in \mathbf{N}^2$. Therefore the corresponding Δ-σ-dimension polynomial is as follows: $\Phi_\eta(t_1, t_2) = t_1 + 2t_2 + 1$. Note that this polynomial expresses the strength of the difference-differential equation $\frac{dy(x)}{dx} - y(x+h) - a(x) = 0$ where $y(x)$ is an unknown function and $a(x)$ belongs to a functional Δ-σ-field K with the derivation $\frac{d}{dx}$ and automorphism $f(x) \mapsto f(x+h)$ where h is a constant of the field (say, if K is a field of functions of a real variable, then h is a real number).

Example 5.12 Let us find the Δ-σ-dimension polynomial that expresses the strength of the difference-differential equation

$$\frac{\partial^2 y(x_1, x_2)}{\partial x_1^2} + \frac{\partial^2 y(x_1, x_2)}{\partial x_2^2} + y(x_1 + h) + a(x) = 0 \tag{5. 6}$$

over some Δ-σ-field of functions of two real variables K, where the basic set of derivations $\Delta = \{\delta_1 = \frac{\partial}{\partial x_1}, \delta_2 = \frac{\partial}{\partial x_2}\}$ has the partition $\Delta = \{\delta_1\} \bigcup \{\delta_1\}$ and σ consists of one automorphisms $\alpha : f(x_1, x_2) \mapsto f(x_1 + h, x_2)\}$ (α is the shift of the first argument of a function by a real number h).

In this case, the associated Δ-σ-extension $K\langle\eta\rangle/K$ is Δ-σ-isomorphic to the field of fractions of the integral Δ-σ-domain $K\{y\}/[\alpha y + \delta_1^2 y + \delta_1^2 y + a]$ (the element $a \in K$ corresponds to the function $a(x)$). Applying Proposition 5.10 we obtain that the characteristic set of the defining ideal of the corresponding Δ-σ-extension $K\langle\eta\rangle/K$ consists of the Δ-σ-polynomials $g_1 = \alpha y + \delta_1^2 y + \delta_1^2 y + a$ and $g_2 = \alpha^{-1} g_1 = \alpha^{-1}\delta_1^2 y + \alpha^{-1}\delta_2^2 y + y + \alpha^{-1}(a)$. With the notation of the proof of Theorem 3.1, the application of the procedure described in this proof, Theorem 4.3(iii), and formula (4.4) leads to the following expressions for the numbers of elements of the sets $U^{(1)}_{r_1 r_2 r_3}$ and $U^{(2)}_{r_1 r_2 r_3}$: $Card\, U^{(1)}_{r_1 r_2 r_3} = r_1 r_2 + 2r_2 r_3 + r_1 + r_2 + 2r_3 + 1$ and $Card\, U^{(2)}_{r_1 r_2 r_3} = 4r_1 r_3 + 2r_2 r_3 - 2r_3$ for all sufficiently large $(r_1, r_2, r_3) \in \mathbf{N}^3$. Thus, the strength of equation (5.6) corresponding to the given partition of the basic set of derivations is expressed by the Δ-σ-polynomial $\Phi_\eta(t_1, t_2, t_3) = t_1 t_2 + 4t_1 t_3 + 4t_2 t_3 + t_1 + t_2 + 1$.

6 Generalized Gröbner bases in free difference-differential modules and difference-differential dimension polynomials

Let K be a difference-differential field of zero characteristic with basic sets $\Delta = \{\delta_1, \dots, \delta_m\}$ and $\sigma = \{\alpha_1, \dots, \alpha_n\}$ of derivations and automorphisms, respectively. Suppose that partition (3.1) of the set of derivations is fixed:
$\Delta = \Delta_1 \bigcup \dots \bigcup \Delta_p \quad (p \geq 1)$ where $\Delta_1 = \{\delta_1, \dots, \delta_{m_1}\}$, $\Delta_2 = \{\delta_{m_1+1}, \dots, \delta_{m_1+m_2}\}, \dots, \Delta_p = \{\delta_{m_1+\dots+m_{p-1}+1}, \dots, \delta_m\}$ $(m_1 + \dots + m_p = m)$.

As before, let Λ denote the free commutative semigroup of all power products of the form $\lambda = \delta_1^{k_1} \dots \delta_m^{k_m} \alpha_1^{l_1} \dots \alpha_n^{l_n}$ where $k_i \in \mathbf{N}$, $l_j \in \mathbf{Z}$ $(1 \leq i \leq m, 1 \leq j \leq n)$ and let $ord_i \lambda$ $(1 \leq i \leq p)$ and $ord_\sigma \lambda$ denote the orders of such an element λ relative to Δ_i and σ, respectively, introduced at the beginning of section 3.

In what follows an expression of the form $\sum_{\lambda \in \Lambda} a_\lambda \lambda$, where $a_\lambda \in K$ for all $\lambda \in \Lambda$ and only finitely many coefficients a_λ are different from zero, is called a difference-differential (or Δ-σ-) operator over K. Two Δ-σ-operators $\sum_{\lambda \in \Lambda} a_\lambda \lambda$ and $\sum_{\lambda \in \Lambda} b_\lambda \lambda$ are considered to be equal if and only if $a_\lambda = b_\lambda$ for all $\lambda \in \Lambda$.

The set of all Δ-σ-operators over K can be equipped with a ring structure if one sets $\sum_{\lambda \in \Lambda} a_\lambda \lambda + \sum_{\lambda \in \Lambda} b_\lambda \lambda = \sum_{\lambda \in \Lambda} (a_\lambda + b_\lambda)\lambda$, $a(\sum_{\lambda \in \Lambda} a_\lambda \lambda) = \sum_{\lambda \in \Lambda} (a a_\lambda)\lambda$, $(\sum_{\lambda \in \Lambda} a_\lambda \lambda)\mu = \sum_{\lambda \in \Lambda} a_\lambda (\lambda\mu)$, $\mu a = \mu(a)\mu$ for any Δ-σ-operators $\sum_{\lambda \in \Lambda} a_\lambda \lambda$, $\sum_{\lambda \in \Lambda} b_\lambda \lambda$ and for any elements $a \in K, \mu \in \Lambda$, and extend these rules by the distributivity. This ring is called the ring of difference-differential (or Δ-σ-) operators over K, it will be denoted by D (or D_K, if the ring K should be specified).

If $u = \sum_{\lambda \in \Lambda} a_\lambda \lambda$ is a Δ-σ-operator over K, then the orders of u relative to the sets Δ_i $(1 \leq i \leq p)$ and σ are defined as numbers $ord_i u =$

$max\{ord_i\lambda | a_\lambda \neq 0\}$ and $ord_\sigma u = max\{ord_\sigma \lambda | a_\lambda \neq 0\}$, respectively. The number $ord\, u = max\{ord\, \lambda | a_\lambda \neq 0\}$ is said to be the order of the Δ-σ-operator u.

For any $r_1, \dots, r_{p+1} \in \mathbf{N}$, let $D_{r_1 \dots r_{p+1}}$ denote the vector K-subspace of D generated by $\Lambda(r_1, \dots, r_{p+1})$. Setting $D_{r_1 \dots r_{p+1}} = 0$ for $(r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1} \setminus \mathbf{N}^{p+1}$, we obtain a family $\{D_{r_1 \dots r_{p+1}} | (r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1}\}$ called a *standard* $(p+1)$*-dimensional filtration* of D.

Obviously, $\bigcup \{D_{r_1 \dots r_{p+1}} | r_1 \dots r_{p+1} \in \mathbf{Z}\} = D$, $D_{r_1 \dots r_{p+1}} \subseteq D_{r_1 \dots r_{i-1}, r_i + 1, r_{i+1}, \dots r_{p+1}}$ for all i, $1 \leq i \leq, p+1$, and $D_{r_1 \dots r_{p+1}} D_{s_1 \dots s_{p+1}} = D_{r_1 + s_1, \dots, r_{p+1} + s_{p+1}}$ for any $r_i, s_i \in \mathbf{N}$, $1 \leq i \leq p+1$.

A left module over the ring D is called a *difference-differential K-module* or a *Δ-σ-K-module.* In other words, a vector K-space M is called a Δ-σ-K-module, if the elements of the set $\Delta \bigcup \sigma^*$ act on M in such a way that $\beta(x+y) = \beta(x) + \beta(y)$, $\beta(\gamma(x)) = \gamma(\beta(x))$, $\delta(ax) = a\delta(x) + \delta(a)x$, $\tau(ax) = \tau(a)\tau(x)$, and $\tau(\tau^{-1}(x)) = x$ for any $\beta, \gamma \in \Delta \bigcup \sigma^*$, $\delta \in \Delta$, $\tau \in \ \sigma^*$, $a \in K$, and $x \in M$.

Definition 6.1 *If M is a Δ-σ-K-module, then a family $\{M_{r_1 \dots r_{p+1}} | (r_1 \dots r_{p+1}) \in \mathbf{Z}\}$ of vector K-subspaces of the module M is called a $(p+1)$-dimensional filtration of M if the following three conditions hold:*

(i) For any fixed integers $r_1, \dots, r_{i-1}$, $r_{i+1}, \dots, r_{p+1}$ $(1 \leq i \leq p+1)$, $M_{r_1 \dots r_i \dots r_{p+1}} \subseteq M_{r_1 \dots r_{i-1}, r_i+1, r_{i+1} \dots r_{p+1}}$ and $M_{r_1 \dots r_{p+1}} = 0$ for all sufficiently small $r_i \in \mathbf{Z}$;

(ii) $\bigcup \{M_{r_1 \dots r_{p+1}} | (r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1}\} = M$;

(iii) $D_{r_1 \dots r_{p+1}} M_{s_1 \dots s_{p+1}} \subseteq M_{r_1 + s_1, \dots, r_{p+1} + s_{p+1}}$ for any $(r_1, \dots, r_{p+1})$, $(s_1, \dots, s_{p+1}) \in \mathbf{Z}^{p+1}$.

If every vector K-space $M_{r_1 \dots r_{p+1}}$ is finitely generated and there exists an element $(h_1, \dots, h_{p+1}) \in \mathbf{Z}^{p+1}$ such that $D_{r_1 \dots r_{p+1}} M_{h_1 \dots h_{p+1}} = M_{r_1 + h_1, \dots, r_{p+1} + h_{p+1}}$ for any
$(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$, then the $(p+1)$-dimensional filtration is called excellent.

It is easy to see that if $u_1, \dots, u_q$ is a finite system of generators of a left D-module M, then the filtration $\{\sum_{i=1}^{q} D_{r_1 \dots r_{p+1}} u_i | \ (r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1}\}$ is excellent.

Let M and N be two Δ-σ-K-modules. A homomorphism of of vector K-spaces $f : M \longrightarrow N$ is called a *Δ-σ-homomorphism* (or *difference-differential homomorphism*), if $f(\tau x) = \tau f(x)$ for any $x \in M, \beta \in \Delta \bigcup \sigma^*$. Surjective (respectively, injective or bijective) Δ-σ-homomorphism is called a Δ-σ-epimorphism (Δ-σ-monomorphism or Δ-σ-isomorphism, respectively).

Let F be a finitely generated free left D-module with free generators $f_1, \dots, f_q$. (Using the "difference-differential" terminology, we also say that F is a free difference-differential K-module (or a free Δ-σ-K-module) with the set of free Δ-σ-generators $\{f_1, \dots, f_q\}$.)

Then F can be considered as a vector K-space with the basis $\{\lambda f_i \,|\, \lambda \in \Lambda$, $1 \leq i \leq q\})$. This set will be denoted by Λf and its elements will be called *terms*. (The fact that the same name is used in section 5 for certain elements of the ring of Δ-σ-polynomials will not cause any confusions.) For any term λf_i, the element λ will be called the *head* of the term.

Since Λf is a basis of F over K, every element $f \in F$ has a unique representation as a linear combination of terms:

$$f = a_1\lambda_1 f_{i_1} + \cdots + a_d\lambda_1 f_{i_d} \qquad (6.\ 1)$$

for some nonzero elements $a_j \in K$ and some $\lambda_j \in \Lambda$ $(1 \leq j \leq d)$. We say that the element f contains a term $\lambda_k f_k$, if this term appears in the representation (6.1) with nonzero coefficient.

We define the orders of a term $\lambda f_j \in \Lambda f$ relative to the sets Δ_i and σ as the corresponding orders of λ: $ord_i(\lambda f_j) = ord_i\lambda$ $(1 \leq i \leq p)$ and $ord_\sigma(\lambda f_j) = ord_\sigma\lambda$. The number $ord\,(\lambda f_j) = ord\,\lambda$ is said to be the order of the term λf_j.

We say that two terms $u = \lambda f_i$ and $v = \lambda' f_j$ are *similar* and write $u \sim v$ if $\lambda \sim \lambda'$. If $u = \lambda e_i$ is a term and $\lambda' \in \Lambda$, we say that u is similar to λ' and write $u \sim \lambda'$ if $\lambda \sim \lambda'$. Furthermore, if $u, v \in \Lambda f$, we say that u *divides* v or v *is a multiple of* u, if $u = \lambda e_i$, $v = \lambda' e_i$ for some e_i and $\lambda|\lambda'$ (in particular, it means that $\lambda \sim \lambda'$).

In what follows we will consider $p+1$ orders $<_1, \dots, <_p, <_\sigma$ on the set Λf defined in the same way as the corresponding orders of the terms in the ring of Δ-σ-polynomials: $\lambda f_j <_i (\text{or } <_\sigma)\ \lambda' f_k$ $(\lambda, \lambda' \in \Lambda,\ 1 \leq j, k \leq q)$ if and only if $\lambda <_i$ (respectively, $<_\sigma$) λ' in Λ, or $\lambda = \lambda'$ and $j < k$.

If an element $f \in F$ is written in the form (6.1) then the greatest element of the set $\{\lambda_1 f_{i_1}, \dots, \lambda_d f_{i_d}\}$ relative to the orders $<_i$ $(1 \leq i \leq p)$ and $<_\sigma$, are called the Δ_i-leader and σ-leader of f; they are denoted by $u_f^{(i)}$ and v_f, respectively.

Definition 6.2 *Let f and g be two elements of the free Δ-σ-K-module F. The element f is said to be reduced with respect to g if f does not contain any multiple λv_g ($\lambda \in \Lambda$) of the σ-leader v_g such that $ord_i(\lambda u_g) \leq ord_i u_f$ for $i = 1, \dots p$. An element $h \in E$ is said to be reduced with respect to a set $\Sigma \subseteq E$, if h is reduced with respect to every element of the set Σ.*

Let us consider p new symbols $z_1, \dots, z_p$, the free commutative semigroup Γ of all power products $\lambda z_1^{k_1} \dots z_p^{k_p}$ $(\lambda \in \Lambda,\ k_i \in \mathbf{N}$ for $i = 1, \dots, p)$, and the set Γf of all elements of the form $\{\gamma f_j | \gamma \in \Gamma, 1 \leq j \leq q\} = \Gamma \times \{e_1, \dots, e_q\}$. If $a = \lambda z_1^{k_1} \dots z_p^{k_p} f_\mu, b = \lambda' z_1^{l_1} \dots z_p^{l_p} f_\nu \in \Gamma f$ $(1 \leq \mu, \nu \leq q)$, we say that a divides b (or b is a multiple of a) and write $a|b$ if $\mu = \nu$, $\lambda|\lambda'$ in Λ, and $k_i \leq l_i$ for $i = 1, \dots, p$.

For any $i = 1, \dots, p$ and for any $f \in F$, let $d(f) = ord_i u_f^{(i)} - ord_i v_f$, and let a mapping $\rho : F \to \Gamma f$ be defined by $\rho(f) = z_1^{d_1(f)} \dots z_p^{d_p(f)} v_f$.

Definition 6.3 *With the above notation, let N be a D-submodule of F. A finite set $G = \{g_1, \dots, g_r\} \subseteq N$ is called a Δ-σ-Gröbner basis of N (with respect to the given partition (3.1) of the basic set Δ) if for any $f \in N$, there exists $g_i \in G$ such that $\rho(g_i)|\rho(f)$.*

Definition 6.4 *Given $f, g, h \in F$, with $g \neq 0$, we say that the element f reduces to h modulo g in one step and write $f \xrightarrow{g} h$ if and only if f contains some term w with a coefficient a such that $v_g | w$, $w = \lambda v_g$ ($\lambda \in \Lambda$, $\lambda \sim v_g$),*

$$h = f - a\,(\lambda_\sigma(lc_\sigma(g)))^{-1}\,\lambda g,$$

and $ord_i(\lambda u_g^{(i)}) \leq ord_i u_f^{(i)}$ for $i = 1, \dots, p$.

Definition 6.5 *Let $f, h \in F$ and let $G = \{g_1, \dots, g_r\}$ be a finite set of nonzero elements of F. We say that f reduces to h modulo G and write $f \xrightarrow{G} h$ if and only if there exists a sequence of elements $g^{(1)}, g^{(2)}, \dots g^{(q)} \in G$ and a sequence of elements $h_1, \dots, h_{q-1} \in F$ such that $f \xrightarrow{g^{(1)}} h_1 \xrightarrow{g^{(2)}} \dots \xrightarrow{g^{(q-1)}} h_{q-1} \xrightarrow{g^{(q)}} h$.*

The introduced notion of reduction is similar to the concept of $(<_k, <_{i_1}, \cdots <_{i_l})$-reduction defined in [9, Section 3]. The following two propositions can be proved in the same way as Theorem 3.6 and Proposition 3.7 of [9].

Proposition 6.6 *With the above notation, let $G = \{g_1, \dots, g_r\} \subseteq F$ be a Δ-σ-Gröbner basis of a D-submodule N of F and let $f \in N$. Then there exist elements $g \in N$ and $Q_1, \dots, Q_r \in D$ such that $f - g = \sum_{i=1}^{r} Q_i g_i$ and g is reduced with respect to G.*

Now we can propose an approach to the computation of the dimension polynomial of a finitely generated Δ-σ-field extension $L = K\langle \eta_1, \dots, \eta_s \rangle$, which is based on the consideration of the corresponding module of Kähler differentials $\Omega_{L|K}$. As it is shown in [3] and [6, Proposition 3.4.46], the L-vector space $\Omega_{L|K}$ can be equipped with the structure of a left module over the ring of Δ-σ-operators $\mathcal{D} = D_L$ over L in such a way that the corresponding action of the elements of $\Delta \bigcup \sigma$ on $\Omega_{L|K}$ satisfy the conditions $\delta(d\zeta) = d\delta(\zeta)$ and $\alpha(d\zeta) = d\alpha(\zeta)$ for any $\zeta \in L$, $\delta \in \Delta$, $\alpha \in \sigma$.

Let us consider $\mathcal{D}$ as a filtered ring with a natural $(p+1)$-dimensional filtration $\{\mathcal{D}_{r_1 \dots r_{p+1}} | (r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1}\}$ where $\mathcal{D}_{r_1 \dots r_{p+1}}$ is the vector L-subspace of $\mathcal{D}$ generated by $\Lambda(r_1, \dots, r_{p+1})$ if all $r_i \geq 0$, and 0 otherwise. Then the finitely generated $\mathcal{D}$-module $M = \Omega_{L|K} = \sum_{i=1}^{s} \mathcal{D} d\eta_i$ has a natural $(p+1)$-dimensional filtration $\{M_{r_1 \dots r_{p+1}} | (r_1, \dots, r_{p+1}) \in \mathbf{Z}^{p+1}\}$ where $M_{r_1 \dots r_{p+1}}$ ($r_i \geq 0$) is the vector L-subspace of $\Omega_{L|K}$ generated by the set $\{d\eta | \eta \in K(\{\lambda\eta_j \,|\, \lambda \in \Lambda(r_1, \dots, r_{p+1}), 1 \leq j \leq s\})$ and $M_{r_1 \dots r_{p+1}} = 0$ whenever at least one r_i is negative. Clearly, this is an excellent filtration of M and, as it is shown in [3] and [6, Theorem 6.4.11],

$$trdeg_K K(\bigcup_{j=1}^{s} \Lambda(r_1, \dots, r_{p+1})\eta_j) = dim_L(M_{r_1 \dots r_{p+1}})$$

for all $(r_1 \dots r_{p+1}) \in \mathbf{N}^{p+1}$.

It follows that the computation of the difference-differential dimension polynomial of the extension L/K can be reduced to the computation of the dimension polynomial associated with the filtration $\{M_{r_1 \dots r_{p+1}}\}$; the existence of such a polynomial is established by the following theorem whose proof is similar to the proof of Theorem 4.2 of [9].

Theorem 6.7 *Let* $\{M_{r_1\dots r_{p+1}} = \sum_{i=1}^{s} \mathcal{D}_{r_1\dots r_p} f_i\}$ *be an excellent* $(p+1)$*-dimensional filtration of a left* $\mathcal{D}$*-module* $M = \sum_{i=1}^{s} \mathcal{D} f_i$ $(f_1, \dots, f_s$ *generate* M *as a* $\mathcal{D}$*-module). Then there exists a polynomial* $\phi(t_1, \dots, t_{p+1}) \in \mathbf{Q}[t_1, \dots, t_{p+1}]$ *such that*

(i) $\phi(r_1, \dots, r_{p+1}) = dim_L M_{r_1\dots r_{p+1}}$ *for all sufficiently large* $(r_1, \dots, r_{p+1}) \in \mathbf{Z}^p$*.*

(ii) $deg_{t_i}\phi \le m_i$ $(1 \le i \le p)$ *and* $deg_{t_{p+1}}\phi \le n$*, so that* $deg\,\phi \le m + n$ *and* $\phi(t_1, \dots, t_{p+1})$ *can be represented as*

$$\phi = \sum_{i_1=0}^{m_1} \cdots \sum_{i_p=0}^{m_p} \sum_{i_{p+1}=0}^{n} a_{i_1\dots i_{p+1}} \binom{t_1 + i_1}{i_1} \dots \binom{t_{p+1} + i_{p+1}}{i_{p+1}}$$

where $a_{i_1\dots i_{p+1}} \in \mathbf{Z}$ *and* $2^n \mid a_{m_1\dots m_p n}$*.*

(iii) *Let* $A = \{(i_1, \dots, i_{p+1}) \in \mathbf{N}^{p+1} \mid 0 \le i_k \le m_k \ (1 \le k \le p),\ 0 \le i_{p+1} \le n$ *and* $a_{i_1\dots i_{p+1}} \ne 0\}$*. Then* $d = deg\,\phi$*,* $a_{m_1\dots m_p n}$*, elements* $(k_1, \dots, k_{p+1}) \in A'$*, the corresponding coefficients* $a_{k_1\dots k_{p+1}}$ *and the coefficients of the terms of total degree* d *do not depend on the choice of the excellent filtration. Also,* $\dfrac{a_{m_1\dots m_p n}}{2^n}$ *is equal to the maximal number of elements of* M *linearly independent over* $\mathcal{D}$*.*

(iv) *Let* E *be a free left* $\mathcal{D}$*-module with free generators* $e_1, \dots, e_s$*. Let* $\pi : E \longrightarrow M$ *be the natural* $\mathcal{D}$*-epimorphism* $(\pi(e_i) = f_i$ *for* $i = 1, \dots, s)$*,* $N = Ker\,\pi$*, and* $G = \{g_1, \dots, g_d\}$ *a* Δ*-*σ*-Gröbner basis of* N*. Furthermore, for any* $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$*, let* $U^{(1)}_{r_1\dots r_{p+1}} = \{u = \lambda e_k \mid ord_i\lambda \le r_i$ *for* $i = 1, \dots, p$*,* $ord_\sigma u \le r_{p+1}$*, and* u *is not a multiple of any* v_{g_j}*,* $j = 1, \dots, d$ $(1 \le k \le d)\}$ *and let* $U^{(2)}_{r_1\dots r_{p+1}} = \{u = \lambda e_k \mid ord_i u \le r_i$ *for* $i = 1, \dots, p$*,* $ord_\sigma u \le r_{p+1}$*, and for every* $\lambda \in \Lambda, g \in G$ *such that* $u = \lambda u_g^{(1)}$*,* $\lambda \sim u_g^{(1)}$*, there exists* $i \in \{1, \dots, p\}$ *such that* $ord_i \lambda u_g^{(i)} > r_i\}$*.*

Then for any $(r_1, \dots, r_{p+1}) \in \mathbf{N}^{p+1}$*, the set* $\pi(U^{(1)}_{r_1\dots r_{p+1}} \bigcup U^{(2)}_{r_1\dots r_{p+1}})$ *is a basis of* $M_{r_1\dots r_{p+1}}$ *over* K*. Therefore* $\phi(r_1, \dots, r_{p+1}) = Card\left(U^{(1)}_{r_1\dots r_{p+1}} \bigcup U^{(2)}_{r_1\dots r_{p+1}}\right)$ *for all sufficiently large* $(r_1, \dots, r_{p+1}) \in \mathbf{Z}^p$*.*

The last theorem shows that the polynomial $\phi(t_1, \dots, t_{p+1})$ can be computed via construction of a Δ-σ-Gröbner basis of the module N mentioned in the last theorem. Such a basis can be obtained with the use of the generalized Buchberger algorithm described in [9] (where $\le_\sigma$ is to play the role of $\le_1$ in the theory developed in [9, Section 3]). Another possible way to construct a Δ-σ-Gröbner basis is to use the relative Gröbner basis technique developed in [11] and [1].

The following example deals with a differential field extension ($\sigma = \emptyset$). It illustrates the method of computation of the multivariate dimension polynomial with the use of the last part of Theorem 6.7 and also shows that multivariate dimension polynomials can carry more invariants of a finitely generated differential (Δ-) field extension than the univariate dimension polynomial considered by E. Kolchin.

Example 6.8 Let K be a differential field with a basic set $\Delta = \{\delta_1, \delta_2, \delta_3\}$ and let $L = K\langle\eta\rangle$ be a Δ-field extension with the defining equation

$$\delta_1^{a+c}\delta_2^{b}\eta + \delta_2^{a+b}\delta_3^{c}\eta + \delta_1^{a}\delta_3^{b+c}\eta = 0$$

where a, b, and c are positive integers. (It means that the defining Δ-ideal of η over K is the Δ-ideal of $K\{y\}$ generated by the Δ-polynomial $f = \delta_1^{a+c}\delta_2^{b}y + \delta_2^{a+b}\delta_3^{c}y + \delta_1^{a}\delta_3^{b+c}y$. Since f is linear, the Δ-ideal $[f]$ is prime.)

As it is shown in [6, Proposition 6.5.5], the kernel N of the natural epimorphism of the free D_L-module $E = D_Le$ onto the module of Kähler differentials $\Omega_{L|K} = D_Ld\eta$ is generated by the element $g = (\delta_1^{a+c}\delta_2^{b} + \delta_2^{a+b}\delta_3^{c} + \delta_1^{a}\delta_3^{b+c})e$, which, obviously, forms a Δ-Gröbner basis of N.

Let $\Phi_\eta(t_1, t_2, t_3)$ be the dimension polynomial associated with the partition

$$\Delta = \{\delta_1\}\bigcup\{\delta_2\}\bigcup\{\delta_3\}.$$

Applying the last part of Theorem 6.7 and the method of evaluation of the sets $U^{(1)}_{r_1,r_2,r_3}$ and $U^{(2)}_{r_1,r_2,r_3}$ described in the proof of Theorem 3.1, we obtain that

$$\Phi_\eta = \left[\binom{t_1+1}{1}\binom{t_2+1}{1}\binom{t_3+1}{1} - \binom{t_1+1-(a+c)}{1}\binom{t_2+1-b}{1}\binom{t_3+1}{1}\right] +$$
$$[(t_1-(a+c)+1)a(t_3+1)+(t_1-(a+c)+1)(t_2-b+1)(b+c)-(t_1-(a+c)+1)a(b+c)] =$$
$$(b+c)t_1t_2 + (a+b)t_1t_3 + (a+c)t_2t_3 + \text{ terms of total degree at most 1.}$$

Note that the corresponding Kolchin differential dimension polynomial, which describes $tr.deg_K K(\Lambda(r)\eta)$, is as follows:

$$\phi_{\eta|K}(t) = \binom{t+3}{3} - \binom{t+3-(a+b+c)}{3} = \frac{a+b+c}{2}t^2 + \text{ terms of degree at most 1.}$$

(One can easily obtain this expression either with the use of the classical Gröbner basis method or by constructing a free resolution for $\Omega_{L|K}$; the corresponding methods and examples can be found, for example, in [6, Chapter 9].)

In this case the Kolchin polynomial $\phi_{\eta|K}(t)$ carries just one differential birational invariant $a+b+c$ while $\Phi_\eta(t_1, t_2, t_3)$ determines three such invariants, $a+b$, $a+c$, and $b+c$, that is, Φ_η determines all three parameters of the defining equation while $\phi_\eta(t)$ gives just the sum of these parameters.

7 Conclusion

We have proved the existence and presented a method of computation of multivariate dimension polynomials associated with difference-differential field extensions and systems of algebraic difference-differential equations. The proposed method, which generalizes the Ritt-Kolchin characteristic set technique, allows one to evaluate the strength of a system of algebraic difference-differential equations in the sense of A. Einstein. We have also presented a method of computation of multivariate difference-differential polynomials based on a generalized Gröbner basis technique that involves several term orderings. Finally, we have found new invariants of a finitely generated difference-differential field extension that are not carried by univariate Kolchin-type dimension polynomials.

References

[1] Dönch, C.; Winkler, F. Bivariate Difference-differential Dimension Polynomials and Their Computation in Maple. *Proceedings of the 8th International Conference on Applied Informatics*, Eger, Hungary, 2010, 211-218.

[2] Einstein, A. *The Meaning of Relativity. Appendix II (Generalization of gravitation theory)*, 4th edn. Princeton, 1953, 133 - 165.

[3] Johnson, Joseph L. Differential dimension polynomials and a fundamental theorem on differential modules. *Amer. J. Math*, 91 (1969), no.1, 239-248.

[4] Kolchin, E. R. The notion of dimension in the theory of algebraic differential equations. *Bull. Amer. Math. Soc.* **70** (1964), 570 - 573.

[5] Kolchin, E. R. *Differential Algebra and Algebraic Groups*. New York: Academic Press, 1973.

[6] Kondrateva, M. V., Levin, A. B., Mikhalev, A. V., Pankratev, E. V. *Differential and Difference Dimension Polynomials*. Dordrecht: Kluwer Academic Publishers, 1999.

[7] Levin, A. B. Characteristic Polynomials of Filtered Difference Modules and Difference Field Extensions.Uspekhi Math. Nauk, Vol. 33 (1978), no. 3, 177 - 178. (In Russian.) English translation: Russian Mathematical Surveys, Vol. 33, no. 3, 165 - 166.

[8] Levin, A. B. Reduced Grobner bases, free difference-differential modules and difference-differential dimension polynomials. *J. Symbolic Comput.*, 29 (2000), 1 - 26.

[9] Levin, A. B. Gröbner bases with respect to several orderings and multivariable dimension polynomials. *J.Symb. Comput.*, 42 (2007), no. 5, 561-578.

[Levin(2008)] Levin, A. B. Difference Algebra. Springer, New York, 2008.

[10] Mikhalev, A. V., Pankratev, E. V. Differential dimension polynomial of a system of differential equations. *Algebra. Collection of papers.* Moscow State Univ. Press, Moscow, 1980, 57 - 67. In Russian.

[11] Zhou, Meng; Winkler, Franz. Computing difference-differential dimension polynomials by relative Gröbner bases in difference-differential modules. *J. Symb. Comput.* 43 (2008), no. 10, 726-745.